

3.5 Biological Resources—Aquatic

This section describes aquatic biological resources—specifically, sensitive fish habitat and important fish species—that could be affected by the proposed program. This section is composed of the following subsections:

- Section 3.5.1, “Environmental Setting,” describes the physical conditions in the study area as they apply to aquatic biological resources.
- Section 3.5.2, “Regulatory Setting,” summarizes federal, State, and regional and local laws and regulations pertinent to evaluation of the proposed program’s impacts on aquatic biological resources.
- Section 3.5.3, “Analysis Methodology and Thresholds of Significance,” describes the methods used to assess the environmental effects of the proposed program and lists the thresholds used to determine the significance of those effects.
- Section 3.5.4, “Environmental Impacts and Mitigation Measures for NTMAs,” discusses the environmental effects of near-term management activities (NTMAs) and identifies mitigation measures for significant environmental effects.
- Section 3.5.5, “Environmental Impacts, Mitigation Measures, and Mitigation Strategies for LTMA’s,” discusses the environmental effects of long-term management activities (LTMA’s), identifies mitigation measures for significant environmental effects, and addresses conditions in which impact analysis would be too speculative for evaluation (CEQA Guidelines, Section 15145).

NTMAs and LTMA’s are described in detail in Section 2.4, “Proposed Management Activities.”

In addition, see Section 3.6, “Biological Resources—Terrestrial,” for a discussion of terrestrial sensitive habitats and sensitive plant and wildlife species.

3.5.1 Environmental Setting

Information Sources Consulted

Sources of information used to prepare this section include the following:

- 1 • *Public Draft Recovery Plan for the Evolutionarily Significant Units of*
2 *Sacramento River Winter-Run Chinook Salmon and Central Valley*
3 *Spring-Run Chinook Salmon and the Distinct Population Segment of*
4 *Central Valley Steelhead* (NMFS 2009a)
- 5 • Discussions published in the *Federal Register* (FR) regarding the
6 statuses of various aquatic biological resources
- 7 • *Inland Fishes of California* (Moyle 2002)
- 8 • *Working Paper: Habitat Restoration Actions to Double Natural*
9 *Production of Anadromous Fish in the Central Valley of California*
10 (USFWS 1995)
- 11 • *Recovery Plan for the Sacramento–San Joaquin Delta Native Fishes*
12 (USFWS 1996)
- 13 • *Final Biological Opinion and Conference Opinion on the Long-Term*
14 *Central Valley Project and State Water Project Operations Criteria*
15 *and Plan. Endangered Species Act Section 7 Consultation* (NMFS
16 2009b)
- 17 • *Formal Endangered Species Consultation on the Proposed*
18 *Coordinated Operations of the Central Valley Project (CVP) and State*
19 *Water Project (SWP)* (USFWS 2008)

20 **Geographic Areas Discussed**

21 Aquatic biological resources are discussed only for the following
22 geographic area within the study area (see Figure 1-4):

- 23 • Extended systemwide planning area (Extended SPA) divided into the
24 Sacramento and San Joaquin Valley and foothills, and the Sacramento–
25 San Joaquin Delta (Delta) and Suisun Marsh

26 The Extended SPA would experience nearly all of the potential effects of
27 the proposed program on aquatic biological resources because almost all of
28 the aquatic habitat in the study area is encompassed by the Extended SPA.
29 In addition, none of the management activities included in the proposed
30 program would be implemented in the SoCal/coastal CVP/SWP service
31 areas. Also, implementation of the proposed program would not result in
32 substantial or long-term reductions in water deliveries to the SoCal/coastal
33 CVP/SWP service areas (see Section 2.6, “No Near- or Long-Term
34 Reduction in Water or Renewable Electricity Deliveries”). Given these
35 conditions, only negligible to no effects on aquatic resources, particularly
36 native species, are expected in the portions of the Sacramento and San

1 Joaquin Valley watersheds that are beyond the Extended SPA or in the
2 SoCal/coastal CVP/SWP service areas outside of the Sacramento and San
3 Joaquin Valley watersheds. Therefore, those geographic areas are not
4 discussed in detail in this section.

5 This section describes fish habitat first in the Sacramento and San Joaquin
6 Valley and foothills, and then in the Delta and Suisun Marsh. Many of the
7 fish species described are present both in waterways of the Sacramento and
8 San Joaquin Valley and foothills and in the Delta and Suisun Marsh at
9 some point in their life history; therefore, this section includes a single
10 discussion of fish species for the entire Extended SPA.

11 ***Fish Habitat in the Extended Systemwide Planning Area***

12 Numerous waterways in the Extended SPA are inhabited by native,
13 sensitive, and/or recreationally valuable fish species that could be affected
14 by the proposed program. The general characteristics of riverine aquatic
15 habitats in the Extended SPA are described below, in the following order:

- 16 • Riverine aquatic habitats in waterways of the Sacramento and San
17 Joaquin Valley and foothills:
 - 18 – Habitats in waterways below human-made structures or natural
19 barriers that block upstream passage of fish
 - 20 – Habitats in the major floodplain bypasses
 - 21 – Reservoir habitat and conditions upstream from the major dams
- 22 • Aquatic habitats within the Delta and Suisun Marsh

23 Names and locations of various rivers, creeks, reservoirs, and other aquatic
24 features in the Extended SPA are shown in Figures 3.13-1 and 3.13-2 in
25 Section 3.13, “Hydrology.”

26 **Sacramento and San Joaquin Valley and Foothills** The Central
27 Valley’s rivers consist of the Sacramento River and its main tributaries
28 (Feather, Yuba, and American rivers and Cache Creek), the eastside
29 tributaries (Cosumnes, Mokelumne, and Calaveras rivers), and the San
30 Joaquin River and its tributaries (Merced, Tuolumne, and Stanislaus
31 rivers). Fish habitat in these rivers is described below, along with habitat
32 found in the major floodplain bypasses and reservoirs in the Sacramento
33 and San Joaquin river systems.

34 *Sacramento River* The Sacramento River is one of California’s largest and
35 most important aquatic ecosystems and supports numerous fish species.
36 Several key tributaries that feed the Sacramento River—the Feather, Yuba,

1 and American rivers and Cache Creek—are also important to aquatic
2 resources. Most of these tributaries are especially important as spawning
3 and rearing areas for anadromous species (species that spend part of their
4 life cycle in the sea and return to their natal freshwater streams to spawn).

5 The Sacramento River is divided into three segments based on
6 geomorphology: the upper segment, which extends from Keswick Dam to
7 Red Bluff Diversion Dam (RBDD); the middle segment, which extends
8 from RBDD to Colusa; and the lower segment, which extends from Colusa
9 to the Delta. The upper Sacramento River tends to retain a more natural
10 geomorphology, but downstream from RBDD, the river channels are
11 confined by levees. Very different types of fish habitats are found in each
12 segment, supporting either different species or different life stages of fish.

13 The upper Sacramento River typically has cool water temperatures because
14 of regulated releases from Shasta and Keswick dams. In this segment, the
15 river channel is stable and confined with little meander, and largely natural
16 with some human-made portions. Riffle habitat with gravel substrates and
17 deep pool habitats are more abundant than in reaches downstream from
18 RBDD. Despite net losses of gravel that occurred after Shasta Dam was
19 constructed, substrates in much of the upper Sacramento River contain
20 gravel that salmonids need to spawn. This stretch of river provides much of
21 the remaining spawning and rearing habitat of several anadromous fishes
22 listed under the Federal Endangered Species Act (ESA) or the California
23 Endangered Species Act (CESA). Therefore, the upper Sacramento River is
24 one of the most sensitive and important sections of any river in California
25 for native anadromous salmonids and green sturgeon.

26 Downstream from RBDD, the middle Sacramento River functions as a
27 large alluvial river with active meander migration along the valley floor.
28 This portion of the river is classified as meandering, and relatively stable,
29 straight sections alternate with more sinuous, dynamic sections (SRCAF
30 2003). Point bars, islands, high and low terraces, instream woody cover,
31 early successional riparian plant growth, and other evidence of river
32 meander and erosion are common. Channel width varies and aquatic
33 habitats consist of shallow riffles, deep runs, deep pools at bends, glides in
34 straight reaches, and shallow vegetated floodplain areas that are inundated
35 during high flows. This stretch of river is used by rearing and migrating
36 salmonids, with some salmonids spawning in the upper portion. Green
37 sturgeon (*Acipenser medirostris*), Sacramento splittail (*Pogonichthys*
38 *macrolepidotus*), striped bass (*Morone saxatilis*), and numerous other
39 species also inhabit the middle Sacramento River.

40 Downstream from Colusa, the lower Sacramento River changes drastically
41 from a dynamic and active meandering channel to a confined (i.e.,

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 restricted from migration) narrow channel. Setback levees exist along
2 portions of the river upstream from Colusa, but levees become much
3 narrower along the river's edge as the river continues south to the Delta.
4 Surrounding agricultural lands encroach directly on the levees, which have
5 cut off the river from most of its riparian corridor, especially on the east
6 side of the river. Most of the levees along the lower Sacramento River are
7 lined with riprap, which reduces the ability of the levees to contribute
8 erodible substrate, reduces habitat variability, and nearly eliminates the
9 processes that lead to the development of complex shaded riverine aquatic
10 (SRA) habitat preferred by native species, and is further described below.
11 Channel width is fairly uniform, and river bends are essentially nonexistent
12 in the straightened channel confined by levees. Therefore, aquatic habitats
13 are fairly homogenous because depth profiles and substrate composition are
14 fairly uniform throughout the reach.

15 Shaded riverine aquatic habitat is defined as the nearshore aquatic area at
16 the interface between the river and adjacent riparian habitat. Such habitat
17 has two principal attributes: an adjacent bank composed of natural, eroding
18 substrates that support riparian vegetation that either overhangs or
19 protrudes into the water; and water that contains variable amounts of
20 instream woody material (IWM) such as leaves, logs, branches, roots, and
21 detritus and has variable velocities, depths, and flows. The USFWS
22 Mitigation Policy has classified shaded riverine aquatic habitat as Resource
23 Category 1 because substantial amounts of such habitat have been lost
24 along the Sacramento River, primarily from levee construction and
25 installation of rock revetment (Fris and Dehaven 1993). The criterion for
26 designating habitat in Resource Category 1 is identified as habitat that is of
27 high value for evaluation species and is unique and irreplaceable on a
28 national basis or in the ecoregion section that could be affected. The
29 mitigation goal for habitat in Resource Category 1 is "no loss of existing
30 habitat value."

31 *Main Tributaries to the Sacramento River* Main tributaries to the
32 Sacramento River within the Sacramento and San Joaquin Valley and
33 foothills consist of the Feather, Yuba, and American rivers and Cache
34 Creek. Aquatic habitats found in these tributaries are discussed below.

35 *Feather River* Aquatic habitats found in the lower Feather River
36 vary as the river flows downstream from DWR's Oroville Dam facilities to
37 the confluence with the Sacramento River at Verona. The low-flow channel
38 of the lower Feather River is approximately 8 miles long and conveys
39 about 600 cubic feet per second (cfs) under current agreement with the
40 California Department of Fish and Game (DFG). The low-flow channel
41 contains mainly riffles and runs, which provide spawning habitat for most
42 Feather River Chinook salmon (*Oncorhynchus tshawytscha*) and steelhead

1 (*O. mykiss*). Also present in the Feather River’s low-flow channel are a
2 series of remnant gravel pit pools/ponds that connect to the main channel.
3 This stretch of the river is fairly confined by levees or high natural banks as
4 it flows through the city of Oroville. The Feather River Salmon and
5 Steelhead Hatchery has been successful in raising fall-run and spring-run
6 Chinook salmon and Central Valley steelhead. However, this effort has
7 caused hatchery-bred salmon and steelhead to dominate the run returns
8 because ocean conditions and in-river habitat have not been optimum for
9 wild production. The hatchery has transitioned to practices that intend to
10 reduce the effects of the hatchery-raised fish on natural production,
11 particularly for the spring-run.

12 From the downstream end of the low-flow channel, the Feather River is
13 fairly active as it meanders south to Marysville; however, this stretch is
14 bordered by active farmland, which confines the river in an incised channel
15 in certain stretches. Minimum flow requirements are being established for
16 the lower Feather River through the Federal Energy Regulatory
17 Commission’s Oroville Facilities relicensing process (DWR 2006). Green
18 and white sturgeon (*Acipenser transmontanus*), American shad (*Alosa*
19 *sapidissima*), and striped bass can be found in the lower Feather River.

20 *Yuba River* The lower Yuba River (downstream from Englebright
21 Dam) is approximately 25 miles long, with about 16 miles of spawning
22 habitat suitable for Chinook salmon. The upstream section of the lower
23 Yuba River is confined by narrow bedrock cliffs and has minimal gravel
24 input. Downstream from the “narrows,” the river consists primarily of
25 cobble/boulder substrate, much of which derives from historic and active
26 mining activities. The mix of deep pools and riffle and run habitat provides
27 healthy habitat conditions for salmonids. The channel is confined by cobble
28 training walls. Daguerre Point Dam, approximately 11 miles upstream
29 from the confluence with the Feather River, is a sediment retention dam
30 that acts as a barrier for sturgeon and other fish (including striped bass and
31 American shad) that cannot pass over the ladders. Because of its substrate
32 condition, the lower Yuba River meanders, with numerous gravel islands in
33 the channel. It also changes its path after high-flow events. Flow
34 requirements, which vary based on water year type, have been set for the
35 Yuba River by the Yuba River Accord to ensure suitable conditions for
36 Chinook salmon, steelhead, and American shad (SWRCB 2008).

37 The lower Yuba River sustains one of the few remaining natural
38 (nonhatchery) populations of Chinook salmon and steelhead in the Central
39 Valley. Strays from the Feather River Salmon and Steelhead Hatchery and
40 other Central Valley hatcheries can also be found there.

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 *American River* The lower American River (downstream from
2 Nimbus Dam) is approximately 23 miles long, has a fairly low gradient,
3 and has riffle, run, glide, and pool habitats. Multiple dams in the watershed
4 have reduced gravel inputs to the system; however, the lower American
5 River contains large gravel bars and side channels in many locations,
6 forming gravel/cobble islands within the channel. Most of the lower
7 American River is surrounded by the American River Parkway, preserving
8 the surrounding riparian zone. The river channel is confined and incised
9 with tall cliffs and bluffs adjacent to the river.

10 The State Water Resources Control Board (SWRCB) established minimum
11 flow requirements for the lower American River through Water Right
12 Decision 893 in 1958. However, the standards in that water right decision
13 were deemed insufficient to protect fisheries by the SWRCB; the U.S.
14 Department of the Interior, Bureau of Reclamation (Reclamation); and the
15 Water Forum (a group of business and agricultural leaders, citizens’
16 groups, environmentalists, water managers, and local governments in the
17 Sacramento region). Therefore, flow management standards were
18 developed with two coequal goals: (1) reliable and safe water supply; and
19 (2) fish, wildlife, recreation, and aesthetic preservation (Water Forum
20 2004). Chinook salmon, steelhead, American shad, and striped bass spawn
21 in the lower American River.

22 *Cache Creek* Within the Extended SPA, Cache Creek extends
23 from the outlet of Clear Lake to the Yolo Bypass. Flows along the entire
24 length of Cache Creek result from dam releases and diversions. Clear Lake
25 is one of the largest freshwater lakes in California, and has the largest
26 surface area of any freshwater lake in California. The lake’s level is
27 regulated by a dam at the outlet to Cache Creek. Before reaching the
28 Sacramento River, Cache Creek flows into the Cache Creek Settling Basin
29 and through the Yolo Bypass.

30 *Eastside Tributaries* Eastside tributaries within the Sacramento and San
31 Joaquin Valley and foothills consist of the Cosumnes, Mokelumne, and
32 Calaveras rivers. These rivers flow into the Delta. Aquatic habitats found in
33 these tributaries are discussed below.

34 *Cosumnes River* The Cosumnes River and its tributaries have no
35 large dams, but Reclamation operates two small dams on this river as part
36 of the CVP. The Cosumnes River supports a spawning population of
37 Chinook salmon, with historic populations estimated up to 17,000
38 spawners. However, since the 1960s, populations have ranged between 0
39 and 5,000 (USFWS 1995). Declines have been assumed to be caused by
40 reduced river flows; at times the river even goes dry or subsurface
41 preventing or delaying upstream migration. The lower reaches of the

1 Cosumnes River also provide important rearing habitat for Sacramento
2 splittail.

3 *Mokelumne River* Anadromous salmonids spawn and rear in
4 approximately 33.5 miles of the regulated Mokelumne River downstream
5 from East Bay Municipal Utility District's Camanche Dam. The 6 miles
6 immediately downstream from the dam are characterized by large gravels
7 and cobble, which change into a sand, mud, and sandstone substrate, with
8 steeper gradients and riffle habitat. This stretch of river contains nearly all
9 the suitable spawning habitat for Chinook salmon and steelhead in the
10 Mokelumne. Downstream from this spawning habitat, the river is low
11 gradient, characterized by alternating bar-complex and flat-water habitats.

12 Mokelumne Dam, approximately 20 miles downstream from Camanche
13 Dam, blocks access by species such as green sturgeon and striped bass that
14 cannot pass over fish ladders. Below the dam, the river is primarily low-
15 gradient run-pool habitat, with riparian debris in the channel and short
16 sections of low-gradient riffles. During low-flow periods, the river channel
17 may develop shallow riffles, impeding passage for both adult and juvenile
18 salmonids. The banks along most of this river reach have substantial
19 riparian growth.

20 *Calaveras River* Approximately 36–38 miles of the Calaveras
21 River downstream from New Hogan Dam are accessible to anadromous
22 fish. Two pathways to the upper portion of the lower Calaveras River,
23 which are split by Bellota Dam, are available: the old Calaveras River
24 channel (36 miles) and Mormon Slough via the Stockton Diverting Canal
25 (38 miles). The latter is the main migration route used by anadromous
26 salmonids due to higher diverted flows into Mormon Slough. Upstream
27 from Bellota Dam, the Calaveras River is considered excellent salmonid
28 habitat, with cold-water, riparian forests that include riparian vegetation
29 and orchards, and floodplain habitat (Fishery Foundation of California
30 2004).

31 Downstream from Bellota Dam, the Calaveras River may have no flows
32 other than tributary inputs from November through mid-April. In the reach
33 downstream from the Stockton Diverting Canal, the river receives urban
34 runoff from storm outlets, and has patches of native riparian and nonnative
35 herbaceous and woody vegetation along its banks. Some sections in this
36 reach have dense native riparian vegetation; others have dense stands of
37 invasive species such as giant reed (*Arundo donax*) and Himalayan
38 blackberry (*Rubus discolor*) that encroach the river channel and banks.
39 Near its confluence with the San Joaquin River, the lower Calaveras River
40 is a narrow, managed, tidally influenced canal (DWR 2007).

1 *San Joaquin River* The San Joaquin River currently does not
2 support spawning anadromous salmonids; however, with the ongoing
3 implementation of the San Joaquin River Restoration Program (SJRRP),
4 self-sustaining populations of spring-run Chinook salmon and steelhead are
5 expected to be reestablished. The time frame for reintroduction is as
6 follows: (1) a Reintroduction Period between the present and December 31,
7 2019; (2) an Interim Period between January 1, 2020, and December 31,
8 2024; (3) a Growth Population Period between January 1, 2025, and
9 December 31, 2040; and (4) a Long-term Period beyond January 1, 2041.
10 The San Joaquin River provides a migratory corridor for salmonids to its
11 major tributary rivers.

12 With successful implementation of the SJRRP, primary habitat for
13 salmonid spawning and rearing would occur in the 37 miles between Friant
14 Dam and Gravelly Ford. The river in this section is an incised, gravel-
15 bedded channel that has been affected by gravel mining, reduced flows and
16 high water temperatures, and introduced aquatic and riparian species.
17 Downstream sections are meandering, low-gradient channels, often with
18 sparse riparian vegetation, low to no flows, and high water temperatures,
19 and are constrained by levees. The section of the San Joaquin River defined
20 by the SJRRP as Reach 4B extends from the Sand Slough Control Structure
21 to where the flood flows in the bypass system rejoin the mainstem of the
22 San Joaquin River and continue to the confluence of the Eastside Bypass.
23 Reach 4B has been perennially dry for more than 40 years, except when
24 agricultural return flows are put through the channel, leaving standing
25 water in many locations. As a result, the upstream end of the Reach 4B
26 channel is poorly defined with dense vegetation and other fill material. The
27 riparian corridor upstream from the Mariposa Bypass is narrow, but nearly
28 unbroken. The downstream portion of Reach 4B contains wider floodplains
29 than upstream reaches and vast areas of natural vegetation.

30 Downstream from the confluence with the Merced River, the San Joaquin
31 River is a relatively wide (approximately 300-foot-wide) channel with little
32 canopy or overhead vegetation and minimal bank cover. Aquatic habitat in
33 the San Joaquin River is characterized primarily by slow-moving glides
34 and pools, is depositional in nature, and has limited water clarity and
35 habitat diversity. Many of the fish species in the lower San Joaquin River
36 use this lower segment of the river to some degree, even if only as a
37 migratory pathway to and from upstream spawning and rearing areas.

38 *Main Tributaries to the San Joaquin River* Main tributaries to the San
39 Joaquin River within the Sacramento and San Joaquin Valley and foothills
40 consist of the Merced, Tuolumne, and Stanislaus rivers. Aquatic habitats
41 found in these tributaries are discussed below.

1 *Merced River* The 10 miles of the Merced River between Crocker-
2 Huffman Dam—the current barrier for fish passage—and State Route 59
3 are where most Chinook salmon and steelhead spawn. This reach has
4 moderate flow and is confined by piles of dredger tailings and sparse
5 riparian vegetation. Gravel mining pits are common for approximately 20
6 miles. The middle stretch of river (approximately 18 miles) is a low-
7 gradient, meandering, and levee-confined system in a narrow corridor,
8 often isolated from its floodplain. The lower 8 miles of the Merced River
9 are sand-bedded, with the most extensive and continuous stand of native
10 riparian vegetation in the river.

11 *Tuolumne River* The Tuolumne River provides roughly 52 miles of
12 waterway accessible to anadromous salmonids. The lower Tuolumne River
13 has two distinct geomorphic zones: a gravel-bedded zone (upper 28 miles)
14 and a sand-embedded zone (lower 24 miles). The gravel-bedded zone is
15 characterized by gravel beds, a discontinuous riparian corridor, gravel
16 mining pits, and dredger tailings. The sand-embedded zone is more
17 affected by urban runoff and encroachment and is dominated by sandy
18 substrate.

19 *Stanislaus River* Aquatic habitats in the lower Stanislaus River
20 vary longitudinally. The river provides spawning, rearing, and/or migratory
21 habitat for a diverse assemblage of common Central Valley native and
22 nonnative fish species. Aquatic habitats consist of riffles, runs, pools, and
23 glides. Floodplain and associated riparian habitat vary with the presence of
24 levees and encroachment of agricultural and urban uses.

25 *Floodplain Bypasses* Three major floodplain bypasses in the Sacramento
26 River system—the Butte Basin, Sutter Bypass, and Yolo Bypass—provide
27 access to broad, inundated floodplain habitat during most years. These
28 bypasses, located along the mainstem of the Sacramento River, include 10
29 overflow structures: six weirs, three flood relief structures, and an
30 emergency overflow roadway. The San Joaquin River system has fewer
31 developed bypass systems—the Chowchilla and Eastside bypasses. These
32 bypasses are not used extensively by fish, but the SJRRP is anticipated to
33 modify portions of the lower bypass system to facilitate salmon migration.

34 Unlike other Sacramento River and Delta habitats, floodplains and
35 floodplain bypasses are seasonally dewatered (as high flows recede) from
36 late spring through autumn. This prevents introduced fish species from
37 establishing year-round dominance except in perennial water sources
38 (Sommer et al. 2003). Moreover, many native fish are adapted to spawn
39 and rear in winter and early spring during the winter flood pulse (Moyle
40 2002). Introduced fish typically spawn from late spring through summer
41 when most of the floodplain is not available to them.

1 *Butte Basin* The Butte Basin lies east of the Sacramento River,
2 extending from the Butte Slough outfall gates near Meridian to Big Chico
3 Creek near Chico Landing. Flood flows are diverted out of the Sacramento
4 River into the Butte Basin and Sutter Bypass via several designated flow
5 relief structures into overflow areas (i.e., low points along the east side of
6 the river) that allow high flood flows to exit the Sacramento River channel.

7 *Sutter Bypass* The Sutter Bypass is a narrow floodwater bypass
8 that conveys Feather and Sacramento river flood flows from the Butte
9 Basin, and Colusa, Moulton, and Tisdale weirs. The bypass area is an
10 expansive land area in Sutter County used mainly for agriculture. In times
11 of high water (when the river stage exceeds 45.45 feet), Sacramento River
12 water enters the bypass through the Butte Slough outfall and Tisdale Weir,
13 inundating the bypass with as much as 12 feet of water. The Sutter Bypass,
14 in turn, conveys flows to the lower Sacramento River region at the Fremont
15 Weir near the confluence with the Feather River, and then into the
16 Sacramento River and the Yolo Bypass.

17 *Yolo Bypass* The Yolo Bypass is a land area of approximately
18 59,000 acres that conveys Sacramento River floodwaters around
19 Sacramento during times of high runoff. Flows are diverted from the
20 Sacramento River into the bypass when the river stage exceeds 33.5 feet
21 (corresponding to 56,000 cfs at Verona). Fremont Weir controls
22 Sacramento River flood stages at Verona by diverting most floodwaters
23 from the Sacramento River into the Yolo Bypass. The Sacramento Weir,
24 just north of Sacramento, diverts flows into the Yolo Bypass downstream
25 of the Fremont Weir. During large flood events, up to 80 percent of
26 Sacramento River flows are diverted into the bypass. The Yolo Bypass also
27 collects flows from the Cache Creek settling basin, Willow Slough,
28 Knights Landing Ridge Cut, and Putah Creek.

29 *Chowchilla Bypass* The Chowchilla Bypass Bifurcation Structure
30 regulates the flow split between the San Joaquin River and the Chowchilla
31 Bypass. The Chowchilla Bypass extends to the confluence of Ash Slough
32 and is approximately 22 miles long, leveed, and 600–700 feet wide. Sand
33 deposits are removed from the bypass as needed, and vegetation is
34 periodically removed from the channel.

35 *Eastside Bypass* The Eastside Bypass circumvents 32.5 miles of
36 the San Joaquin River. This bypass extends from the confluence of Ash
37 Slough and the Chowchilla Bypass to the confluence with the San Joaquin
38 River. Within the Eastside Bypass, the Mariposa Bypass connects the
39 Eastside Bypass to Reach 4B of the San Joaquin River. The Eastside
40 Bypass is subdivided into three sections:

- 1 • Section 1 extends from Ash Slough to the Sand Slough Bypass
2 confluence. This reach receives flows from the Chowchilla River at
3 river mile (RM) 136.
- 4 • Section 2 extends from the Sand Slough Bypass to the head of the
5 Mariposa Bypass at RM 147.2.
- 6 • Section 3 extends from the head of the Mariposa Bypass to the San
7 Joaquin River at RM 168.5, and receives flows from Deadman, Owens,
8 and Bear creeks.

9 *Reservoirs* Reservoirs have been one of the major sport-fish habitats in the
10 Central Valley since the advent of the SWP and CVP. Numerous reservoirs
11 exist in the Extended SPA: Shasta Lake, Lake Oroville, Englebright
12 Reservoir, Clear Lake, Folsom Lake, Camanche Reservoir, New Hogan
13 Reservoir, New Melones Reservoir, New Don Pedro Reservoir, Lake
14 McClure, and Millerton Lake.

15 The nature of each reservoir and its fish fauna is determined by the
16 reservoir's size, elevation, location, and water quality. In general,
17 reservoirs are much less productive per surface acre than natural lakes
18 because reservoirs are generally deep, steep-sloped basins, and fluctuating
19 water levels greatly limit habitat diversity. Central Valley reservoirs
20 generally fall into one of two categories: warm-water reservoirs, suitable
21 for black bass, sunfish, and catfish; or "two-story" reservoirs that contain a
22 zone of deep, well-oxygenated water in summer that is suitable for trout,
23 topped by warm-water surface waters suitable for black bass, sunfish, and
24 catfish.

25 Warm-water reservoirs usually have low fertility and yield relatively small
26 crops of game fish. Because of extensive drawdowns, inshore zones
27 inhabited by warm-water species are usually relatively unproductive. The
28 deep, open-water portion of a large reservoir also does not provide
29 satisfactory habitat for most game fish.

30 Extreme water-level fluctuation in reservoirs is perhaps the most important
31 environmental factor influencing the productivity of reservoir fish
32 populations, and is a direct result of reservoir management priorities.
33 Fluctuating water levels are largely responsible for other fishery
34 management problems, such as limited cover habitat, limited littoral
35 habitat, and shoreline erosion. Central Valley reservoirs operate to store
36 water during winter and spring and then release water in summer and fall.
37 This pattern of storage and releases results in variable, seasonal availability
38 of water in reservoirs. Surface-water elevation fluctuations in some Central
39 Valley reservoirs could exceed 100 vertical feet annually.

1 **Delta and Suisun Marsh** The Delta, and Suisun Marsh on the western
2 edge of the Delta, are located at the confluence of the Sacramento and San
3 Joaquin rivers. This is the most important, complex, and controversial
4 geographic area in California for anadromous fish production, estuarine
5 fish species, introduced fish species, and distribution of water resources for
6 numerous beneficial uses.

7 The Delta's environmental conditions depend primarily on the physical
8 structure of Delta levees and channels, inflow volume and source, Delta
9 Cross Channel operations, Delta exports and diversions, and tides. The
10 CVP and SWP affect Delta conditions primarily by controlling upstream
11 storage and diversions, reservoir releases, Delta water conveyance
12 pathways, and Delta exports and diversions. These factors also determine
13 Delta outflow and the location of the entrapment zone, an area of high
14 organic carbon that is critically important to numerous fish and invertebrate
15 species and to the overall ecology of the Delta and Suisun Marsh.

16 In addition to these physical factors, environmental conditions contribute to
17 interactive, cumulative conditions that substantially affect Delta fish
18 populations. Water temperature, predation (primarily by introduced fish
19 species), food production and availability, competition with introduced
20 exotic fish and invertebrate species, reduced habitat complexity, and
21 pollutant concentrations are all important contributors to cumulative
22 conditions.

23 An estimated 25 percent of all sportfishing for warm-water and
24 anadromous species and 80 percent of California's commercial fishery
25 depend on species that live in or migrate through the Delta. The Delta
26 serves as a migration corridor for all Central Valley anadromous species as
27 they return to their natal rivers to spawn, and during juvenile outmigration
28 downstream to the ocean. Adult Chinook salmon move through the Delta
29 during most months of the year. Salmon and steelhead juveniles depend on
30 the Delta as transient rearing habitat while they migrate through the system
31 to the ocean; these juveniles could remain for several months, feeding in
32 marshes, tidal flats, and sloughs. All life stages of striped bass are found in
33 the Delta and approximately 45 percent of striped bass spawn there.
34 Numerous resident species live in the Delta year round, such as delta smelt,
35 Sacramento splittail, and introduced threadfin shad (*Dorosoma petenense*).

36 Since about 2002, four pelagic (occupying the open water) fish species
37 have been subject to the Pelagic Organism Decline (POD) (Sommer et al.
38 2007). The POD refers to the sudden, overlapping declines of pelagic fishes
39 in the San Francisco Bay/Sacramento–San Joaquin Delta estuary (Bay-
40 Delta) that were first recognized in data collected in 2002–2004. The
41 species identified in the POD consist of delta smelt, longfin smelt, threadfin

1 shad, and (age-0) striped bass. Together, these species account for most of
2 the resident pelagic fish biomass in the tidal water upstream from X2, the
3 position at which 2 parts per thousand (ppt) salinity occurs in the Bay-
4 Delta.

5 The declines of three of the four POD species became noticeable between
6 2001 and 2002; however, studies have revealed that at least for delta smelt,
7 the POD downtrend actually began earlier, around 1999 (Manly and
8 Chotkowski 2006, as cited in USFWS 2008). Abundance indices for the
9 POD species since 2001 have included record lows for all but threadfin
10 shad. The causes of the POD and earlier declines are not fully understood,
11 but studies are under way to evaluate potential causes. Among these
12 potential causes are the stock-recruitment relationship (i.e., previous
13 abundance), a decrease in habitat carrying capacity or production potential,
14 predation and entrainment, and a decline in primary productivity (Moyle et
15 al. 1992; Bennett 2005; Feyrer et al. 2007; Baxter et al. 2010). In 2011,
16 both delta smelt and longfin smelt populations showed an increase, with
17 delta smelt populations at their highest since 2001 and longfin smelt at their
18 highest since 2006. However, these increases are still a fraction of historic
19 abundances.

20 ***Fish Species in the Extended Systemwide Planning Area***

21 Various fish species in the Extended SPA are considered important either
22 for their legal status or for their economic, ecological, or recreational value.
23 The lower Sacramento and San Joaquin rivers, their tributaries, and the
24 Delta and Suisun Marsh provide vital fish spawning, rearing, and/or
25 migratory habitat for a diverse assemblage of native and nonnative species.
26 The key life stages and needs of the species of primary management
27 concern with the greatest potential to be affected by the proposed program
28 are discussed below. These species collectively represent a diversity of life
29 histories and environmental/habitat requirements, and they are among the
30 most sensitive to environmental perturbation; therefore, findings from
31 assessments of these species can be effectively used to make inferences
32 about other fish species that use the Extended SPA. The seasonal timing of
33 important life stages for these species in the study area is presented
34 (represented by the gray boxes) in Table 3.5-1. Special-status species and
35 occurrence within the Extended SPA are shown in Table 3.5-2.

3.0 Environmental Setting, Impacts, and Mitigation Measures
 3.5 Biological Resources—Aquatic

Table 3.5-1. Life History and Distributions of Life Stages for Key Fish Species in the Extended Systemwide Planning Area

Life Stage/ Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Fall-Run Chinook Salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Late Fall-Run Chinook Salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Winter-Run Chinook Salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Spring-Run Chinook Salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Steelhead												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Green Sturgeon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Pacific Lamprey												
Adult migration												
Spawning												
Larvae and Juvenile Rearing												
Emigration												
Delta Smelt												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Estuarine rearing												
Longfin Smelt												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Estuarine rearing												
Sacramento Splittail												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Adult and juvenile rearing												

Table 3.5-1. Life History and Distributions of Life Stages for Key Fish Species in the Extended Systemwide Planning Area (contd.)

Life Stage/ Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Hardhead												
Adult foraging and spawning												
Spawning												
Larvae and juvenile rearing												
Adult and juvenile rearing												
Striped bass												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Adult and juvenile rearing												

Sources: Vogel and Marine 1991; Moyle 2002; Wang 1986; NMFS 2005

Key:

 = period of potential occurrence

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

Table 3.5-2. Special-Status Species and Occurrence Within the Extended Systemwide Planning Area

Species	Status	Distribution
Central Valley fall-/late fall–run Chinook salmon	SSC, SC	Sacramento and San Joaquin rivers and their major tributaries, Eastside Tributaries; Delta, Suisun Bay; Suisun and Napa marshes, San Francisco Bay, Pacific Ocean
Central Valley spring-run Chinook salmon	ST, FT	Feather, Sacramento, and Yuba rivers; Beegum, Battle, Clear, Cottonwood, Antelope, Mill, Deer, Butte, and Big Chico creeks; Delta, Suisun Bay; Suisun and Napa marshes, San Francisco Bay, Pacific Ocean
Sacramento River winter-run Chinook salmon	SE, FE	Sacramento River (Keswick Dam to Chipps Island); Delta, Suisun Bay; Suisun and Napa marshes, San Francisco Bay, Pacific Ocean
Central Valley steelhead	FT	Sacramento and San Joaquin rivers and their major tributaries, Eastside Tributaries; Delta, Suisun Bay; Suisun and Napa marshes, San Francisco Bay, Pacific Ocean
Southern distinct population segment of the North American green sturgeon	FT	Sacramento River; lower Feather River; Yuba River; ; Delta, Suisun Bay; Suisun and Napa marshes; San Francisco Bay, Pacific Ocean
Delta smelt	SE, FT	Lower Sacramento River; lower San Joaquin River, San Joaquin River ; Delta, San Francisco Bay, Pacific Ocean
Longfin smelt	ST, Federal Status Review underway	Klamath, Eel, and San Francisco Bay/Sacramento–San Joaquin Delta estuaries; Delta, Suisun Bay; Suisun and Napa marshes , San Francisco Bay
Sacramento splittail	SSC	Delta; Suisun Bay; Suisun and Napa marshes; Sacramento and San Joaquin rivers
Hardhead	SSC, SC, USFS sensitive species	Low/mid-elevation streams in Sacramento and San Joaquin river watersheds
Pacific lamprey	No status	Sacramento and San Joaquin rivers and their major tributaries; Delta, San Francisco Bay, Pacific Ocean
River lamprey	SSC	Sacramento and San Joaquin rivers and their major tributaries
Striped bass	No status	Sacramento River and its major tributaries; San Joaquin River; ; Delta, San Francisco Bay, Pacific Ocean

Source: Data compiled by MWH in 2011

Key:

Delta = Sacramento–San Joaquin Delta

SSC = State species of special concern

SC = Federal species of concern

ST = State threatened species

SE = State endangered species

FT = Federal threatened species

FE = Federal endangered species

USFS = U.S. Forest Service

1 **Chinook Salmon** The Sacramento River and its tributaries support four
2 separate runs of Chinook salmon—fall-run, late fall–run, winter-run, and
3 spring-run, denoting when adults enter freshwater and begin their upstream
4 migration. Spring-run Chinook salmon are both State-listed and federally
5 listed as threatened. Winter-run Chinook salmon are both State-listed and
6 federally listed as endangered. Fall- and late fall–run salmon are not
7 currently listed by either the State or federal government, but because of
8 low population numbers, the State identified them as a species of special
9 concern and the National Marine Fisheries Service (NMFS) identified them

1 as a species of concern. Most Chinook salmon and all steelhead spawn
2 between Keswick Dam and RBDD, but fall-run Chinook salmon spawn as
3 far downstream as Colusa. The San Joaquin River tributaries currently
4 support only fall-run Chinook salmon, although with implementation of the
5 SJRRP, spring-run Chinook salmon are expected to be successfully
6 reintroduced.

7 *Fall-Run Chinook Salmon* Fall-run Chinook salmon represent about 80
8 percent of the total Chinook salmon produced in the Sacramento River
9 drainage and 100 percent of the Chinook salmon in the eastside tributaries
10 and San Joaquin River watershed (Kjelson et al. 1982). On March 9, 1998
11 (63 FR 11481), NMFS issued a proposed rule to list fall-run Chinook
12 salmon as threatened; however, NMFS determined that the fall-run did not
13 warrant listing and identified it as a candidate species (64 FR 50393,
14 September 16, 1999). NMFS also determined that both late fall–run and
15 fall-run are a single evolutionarily significant unit (ESU), but because they
16 are separate in timing and effects, they are distinguished as separate in this
17 document. They later designated Central Valley fall- and late fall–run as a
18 species of concern (69 FR 19975, April 15, 2004).

19 Adult fall-run Chinook salmon migrate from July through December. Fall-
20 run Chinook salmon spawn between early October and late December, and
21 incubation takes place from October through March. Spawning activity
22 peaks in October and November as water temperature drops. Fall-run
23 Chinook salmon move upstream from the ocean in the late summer and
24 early fall in mature condition and spawn soon after arriving at their
25 spawning grounds. Juvenile Chinook salmon emerge from the gravel and
26 migrate downstream to the ocean soon after emerging, rearing in the
27 streams for only few months.

28 *Late Fall–Run Chinook Salmon* Late fall–run Chinook salmon mostly
29 inhabit the Sacramento River, spawning upstream from RBDD. They
30 migrate into the Sacramento River between October and April and spawn
31 from January through April. Spawning activity peaks in February and
32 March, followed by egg incubation from January through June and fry
33 emergence from April through June. Rearing and emigration of fry and
34 smolts occur from April through December. Juvenile late fall–run Chinook
35 salmon rear in the streams during the summer; in some streams, they
36 remain throughout the year.

37 *Sacramento River Winter-Run Chinook Salmon* The Sacramento River
38 upstream from RBDD is the only currently known spawning reach for
39 winter-run Chinook salmon, which have been in a major decline since the
40 1960s. The sharp decline in adult escapement during the late 1980s and
41 early 1990s prompted listing of the winter-run Chinook salmon ESU as

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 endangered under the ESA on January 4, 1994 (59 FR 440), and under the
2 CESA on September 22, 1989. In 2009 NMFS submitted a draft recovery
3 plan that includes proposed management actions to help protect
4 Sacramento River winter-run Chinook salmon (NMFS 2009a).

5 The portion of the Sacramento River from Keswick Dam to Chipps Island
6 (near Suisun Bay), all waters in the Delta westward from Chipps Island to
7 the Carquinez Bridge, all waters of San Pablo Bay, and all waters of San
8 Francisco Bay north of the San Francisco–Oakland Bay Bridge have been
9 designated as critical habitat for winter-run Chinook salmon (58 FR 33212,
10 June 16, 1993). Critical habitat consists of the river water, river bottom,
11 and adjacent riparian zone (i.e., those adjacent terrestrial areas that directly
12 affect a freshwater aquatic ecosystem).

13 Adult winter-run Chinook salmon leave the ocean and migrate through the
14 Delta into the Sacramento River from November through July, passing
15 RBDD on the Sacramento River between mid-December and July.
16 Spawning takes place from mid-April through August, and egg incubation
17 continues through mid-October.

18 Juvenile winter-run Chinook salmon rear in the upper Sacramento River
19 from July through March (Hallock and Fisher 1985). Juveniles move
20 downstream in the river from August through October, and possibly
21 through November. Juveniles have been observed in the Delta from
22 October through December. In general, juvenile abundance in the Delta
23 increases in response to increased Sacramento River flow (USFWS 1995).

24 Winter-run Chinook salmon smolts (i.e., juveniles that are physiologically
25 ready to enter seawater) may migrate through the Delta and San Francisco
26 Bay to the ocean from December through May (Stevens 1989). The
27 Sacramento River channel is their main migration route through the Delta.

28 *Spring-Run Chinook Salmon* On September 16, 1999, NMFS listed the
29 Central Valley spring-run Chinook salmon ESU as threatened under the
30 ESA. The Central Valley spring-run Chinook salmon ESU consists of all
31 naturally spawned populations of spring-run Chinook salmon in the
32 Sacramento River and its tributaries, as well as artificially propagated
33 Feather River spring-run Chinook salmon (70 FR 37177, June 28, 2005). In
34 2009 NMFS submitted a draft recovery plan that includes proposed
35 management actions to help protect Central Valley spring-run Chinook
36 salmon (NMFS 2009a). The State listed spring-run as threatened on
37 February 5, 1999.

38 Critical habitat for spring-run Chinook salmon comprises roughly 1,272
39 miles of occupied stream habitat and 427 square miles of estuarine habitat.

1 This critical habitat encompasses the lower Feather River; the Sacramento
2 and Yuba rivers; Beegum, Battle, Clear, Cottonwood, Antelope, Mill, Deer,
3 Butte, and Big Chico creeks; the north Delta (the central and south Delta
4 were excluded); and Suisun, San Pablo, and north San Francisco bays (70
5 FR 52488, September 2, 2005).

6 Adult spring-run Chinook salmon enter the mainstem Sacramento River in
7 February and March, when they are sexually immature. Adults hold over
8 summer in deep, cold-water pools near spawning habitat until they spawn
9 between late August and October. In most locations, juveniles emerge in
10 November and December, but they may emerge later when water
11 temperature is cooler. Spring-run Chinook salmon may migrate
12 downstream as young-of-year juveniles or as yearlings. Based on
13 observations in Butte Creek and the Sacramento River, young-of-year
14 juveniles migrate from November through June. Yearling spring-run
15 Chinook salmon migrate from October through March, with peak migration
16 in November (Cramer and Demko 1997, Hill and Webber 1999).

17 **Central Valley Steelhead** On March 19, 1998, the naturally spawned
18 Central Valley steelhead was federally listed as threatened by NMFS (63
19 FR 13347). The Central Valley steelhead distinct population segment
20 (DPS) includes all naturally spawned populations of steelhead (and their
21 progeny) in the Sacramento and San Joaquin rivers and their tributaries.
22 Resident rainbow trout were previously included as protected fish;
23 however, in 2006, NMFS directed that only the anadromous form should be
24 listed as threatened, and the resident form did not warrant listing (71 FR
25 834, January 5, 2006). It is difficult, if not impossible, to distinguish
26 anadromous and resident juvenile *O. mykiss*, but adults are different
27 enough in appearance to distinguish. In 2009 NMFS submitted a draft
28 recovery plan that includes proposed management actions to help protect
29 Central Valley steelhead (NMFS 2009a).

30 Critical habitat for Central Valley steelhead encompasses the lower Feather
31 River; Battle, Cottonwood, Antelope, Mill, Deer, Big Chico, and Butte
32 creeks; the Sacramento, Yuba, American, Cosumnes, Mokelumne,
33 Calaveras, San Joaquin, Merced, Tuolumne, and Stanislaus rivers; and the
34 Delta.

35 Central Valley steelhead migrate upstream from July through May.
36 Spawning in the Sacramento River basin typically occurs from late
37 December through April, with most spawning occurring from January
38 through March. Unlike Chinook salmon, which die after spawning,
39 steelhead can survive spawning and live to spawn more than once. The
40 eggs hatch 19–80 days after spawning, depending on water temperature
41 (warmer temperatures result in faster hatching times), and the young

1 remain in the gravel for several weeks before emerging as fry (Raleigh et
2 al. 1984).

3 Steelhead juveniles rear a minimum of 1 year, and typically 2 years, in
4 freshwater before emigrating to the ocean as smolts. Smolt emigration
5 generally occurs from November through May, although based on salvage
6 data at State and federal pumping plants in the Delta, March and April
7 appear to be the peak months for emigration in most years. After spending
8 2–3 years in the ocean, steelhead return to their natal streams to spawn as
9 4- or 5-year-olds.

10 **Green Sturgeon** North American green sturgeon have been separated into
11 two DPSs: the northern DPS (all populations north of and including the Eel
12 River) and the southern DPS (coastal and Central Valley populations south
13 of the Eel River). The southern DPS is federally listed as threatened under
14 the ESA (71 FR 17757, April 7, 2006). In the Extended SPA, critical
15 habitat has been designated for the Sacramento River, the lower Feather
16 and Yuba rivers, the Delta, and Suisun, San Pablo, and San Francisco bays
17 (74 FR 52300, October 9, 2009).

18 Little is known about the life history of green sturgeon because of its low
19 abundance, low sportfishing value, and limited spawning distribution;
20 however, the spawning and larval ecology of green sturgeon are assumed to
21 be similar to those of white sturgeon (Moyle 2002; Beamesderfer and
22 Webb 2002). Green sturgeon are mostly marine fish but migrate into rivers
23 to spawn. Green sturgeon also make extensive ocean migrations;
24 consequently, most recoveries of individuals tagged in San Pablo Bay have
25 come from the ocean and from rivers and estuaries in Oregon and
26 Washington.

27 Within estuaries, green sturgeon reportedly tend to concentrate in deep
28 areas with soft bottoms. In rivers, adult (and juvenile) green sturgeon have
29 been observed primarily on clean sand (EPIC et al. 2001). Adult green
30 sturgeon are benthic, usually found in the Sacramento River in deep, off-
31 channel areas with little current. Both the upstream and downstream
32 migrations of green sturgeon begin in late February and continue through
33 July (Beamesderfer and Webb 2002), and reach above RBDD to near
34 Keswick Dam.

35 Most females reach sexual maturity at 20–25 years while males reach
36 sexual maturity at 15–17 years (Beamesderfer and Webb 2002). Green
37 sturgeon are thought to spawn every 3–5 years (70 FR 65, April 6, 2005)
38 from February to July, with a peak in mid-April to mid-June (Kohlhorst
39 1976, Moyle 2002, Beamesderfer and Webb 2002). The reported range of
40 preferred/optimal water temperatures for green sturgeon spawning is

1 unclear, but spawning success is related to water temperature
2 (Beamesderfer and Webb 2002). Green sturgeon spawn in deep pools in
3 large, turbulent rivers (Moyle et al. 1992); the preferred spawning substrate
4 is likely large cobble-containing crevices in which eggs can become
5 trapped and develop, but may also range from clean sand to bedrock (EPIC
6 et al. 2001, Beamesderfer and Webb 2002).

7 Sturgeon eggs have been found in the Sacramento River from mid-
8 February through July (Kohlhorst 1976; Moyle 2002; Beamesderfer and
9 Webb 2002). The importance of water quality is uncertain, but silt is
10 known to prevent green sturgeon eggs from adhering to each other
11 (USFWS 1996), and sand and silt may suffocate the eggs (EPIC et al.
12 2001). Water temperatures above 68 degrees Fahrenheit (°F) (20 degrees
13 Celsius (°C)) are reportedly lethal to green sturgeon embryos
14 (Beamesderfer and Webb 2002).

15 Juvenile green sturgeon reportedly occur in shallow water (Radtke 1966)
16 and probably move to deeper more saline areas as they grow (EPIC et al.
17 2001). Rearing juveniles remain in freshwater for 1–4 years before
18 returning to their marine environment (Beamesderfer and Webb 2002;
19 EPIC et al. 2001). Juveniles in the Delta feed primarily on opossum shrimp
20 and amphipods (Radtke 1966, Moyle 2002).

21 **Delta Smelt** Delta smelt was federally listed as threatened in 1993 (58 FR
22 12854, March 5, 1993); critical habitat was designated on December 19,
23 1994. Critical habitat includes the portion of the Sacramento River from
24 Keswick Dam to Chipps Island, all waters westward from Chipps Island to
25 the Carquinez Bridge, all waters of San Pablo Bay, and all waters of San
26 Francisco Bay north of the San Francisco–Oakland Bay Bridge. Delta smelt
27 were upgraded from threatened to endangered status under the CESA on
28 January 20, 2010.

29 Delta smelt are endemic to the Delta. During the spawning season, adults
30 move into the Delta's channels and sloughs. When Delta outflows are high,
31 delta smelt may occur in San Pablo Bay. Delta smelt have relatively low
32 fecundity and most live for 1 year (Moyle 2002).

33 Estuarine rearing habitat for juvenile and adult delta smelt is typically
34 found in the waters of the lower Delta and Suisun Bay, where salinity is 2–
35 7 ppt. Delta smelt tolerate 0–19 ppt salinity. They typically occupy open
36 shallow waters (less than 10 feet deep), but they also occur in the main
37 channel in the areas where freshwater and brackish water mix.

38 Adult delta smelt begin a spawning migration, which may encompass
39 several months, toward the upper Delta and freshwater in December or

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 January. Spawning occurs between February and July, with peak spawning
2 from April through mid-May. Delta smelt spawn in shallow edge-waters in
3 upstream Delta channels, including the Sacramento River above Rio Vista,
4 Cache Slough, Lindsey Slough, and Barker Slough (near the downstream
5 end of the Yolo Bypass). Eggs are broadcast over the bottom of the
6 channel, where they attach to firm sediment, woody material, and
7 vegetation. Larval smelt feed on rotifers and other zooplankton. Larvae and
8 juveniles gradually move downstream toward rearing habitat in the
9 estuarine mixing zone.

10 **Longfin Smelt** DFG has designated the longfin smelt (*Spirinchus*
11 *thaleichthys*) as threatened and fully protected under the CESA, but the
12 U.S. Fish and Wildlife Service (USFWS) did not pursue listing longfin
13 smelt under the federal ESA. Historically, longfin smelt populations were
14 found in the Klamath, Eel, and Bay-Delta estuaries and in Humboldt Bay.
15 In the Central Valley, longfin smelt are rarely found upstream from Rio
16 Vista or Medford Island (northwest of Stockton) in the Delta. Adults
17 concentrate in Suisun, San Pablo, and north San Francisco bays (Moyle
18 2002).

19 Longfin smelt are found in San Pablo Bay from April through June and
20 disperse in late summer. In fall and winter, yearlings move upstream into
21 freshwater to spawn. Longfin smelt spawn downstream from Medford
22 Island in the San Joaquin River and downstream from Rio Vista on the
23 Sacramento River. Spawning may occur as early as November, and larval
24 surveys indicate that it may extend into June (Moyle 2002).

25 High outflows transport the larvae into Suisun and San Pablo bays. In low-
26 outflow years, larvae move into the western Delta and Suisun Bay. Higher
27 outflows reflect positively in juvenile survival and adult abundance.
28 Rearing habitat is better in Suisun and San Pablo bays because juveniles
29 require brackish water in the 2- to 18-ppt range. If juveniles stay in the
30 Delta, they may become entrained and exposed to more adverse conditions
31 (Moyle 2002), such as continued predation by introduced fish species.

32 **Sacramento Splittail** On September 22, 2003, USFWS delisted the native
33 Sacramento splittail as a threatened species because habitat restoration
34 actions such as the CALFED Bay-Delta Program (CALFED) and the
35 Central Valley Project Improvement Act (CVPIA) were expected to
36 prevent the splittail from becoming endangered in the foreseeable future
37 (68 FR 55139, September 22, 2003). Splittail is identified as a species of
38 special concern under the CESA.

39 Splittail are found primarily in the Delta, Suisun Bay, Suisun Marsh, and
40 Napa Marsh, but juveniles have been found in the Sacramento River as far

1 upstream as RBDD (Sommer et al. 1997), and in the San Joaquin River as
2 far upstream as Salt Slough (just upstream from the Merced River
3 confluence (Moyle 2002). Sommer et al. (1997, 2002) found that the Yolo
4 and Sutter bypasses provide important spawning habitat for splittail.

5 Adult splittail migrate from Suisun Bay and the Delta to upstream
6 spawning and rearing habitat from December through April. This species
7 prefers low water velocities for spawning and early rearing. Splittail spawn
8 in Suisun Marsh in late April and May and in the upper Delta and lower
9 reaches and flood bypasses of the Sacramento and San Joaquin rivers
10 between early March and May (Moyle et al. 1995). Spawning has been
11 observed to occur as early as January and may continue through early July
12 (Wang 1986; Moyle 2002).

13 Larval splittail are commonly found in shallow, vegetated areas near
14 spawning habitat. Larvae eventually move into deeper and more open-
15 water habitat as they grow. During late winter and spring, young-of-year
16 juvenile splittail (i.e., production from spawning in the current year) are
17 found in sloughs, rivers, and Delta channels near spawning habitat.
18 Juvenile splittail gradually move from shallow, nearshore areas to the
19 deeper open-water habitat of Suisun and San Pablo bays (Wang 1986). In
20 areas upstream from the Delta, juvenile splittail can be expected to be
21 present in the flood bypasses when these areas are inundated during winter
22 and spring (Jones & Stokes Associates 1993; Sommer et al. 1997).

23 **Hardhead** Hardhead are widely distributed throughout the low- to mid-
24 elevation streams in the main Sacramento–San Joaquin River drainage.
25 Undisturbed portions of larger streams at low to middle elevations are
26 preferred by hardhead. They are fairly intolerant of low-oxygen waters,
27 particularly at higher water temperatures. Pools with sand-gravel substrates
28 and slow water velocities are the preferred habitat; adult fish inhabit the
29 lower half of the water column, while the juvenile fish remain in the
30 shallow water closer to stream edges. Hardhead typically feed on small
31 invertebrates and aquatic plants at the bottom of quiet water (Moyle 2002).
32 Hardhead, a native species, is a State species of special concern, and a U.S.
33 Forest Service sensitive species in California.

34 **Pacific Lamprey** Pacific lamprey (*Entosphenus tridentata*) are found
35 within the Extended SPA and are native to California waters. Pacific
36 lamprey enter freshwater anywhere from a few months to a few years prior
37 to spawning during the late spring through the fall, indicating there are
38 potentially multiple runs. Spawning occurs in low gradient reaches, in
39 gravel typically in pool tails or riffles. After spawning, the ammocoetes
40 (larvae) spend 3 to 4 years in freshwater, feeding on detritus, diatoms and
41 algae, before metamorphosing into juveniles. Ammocoetes rear in sand and

1 mud substrates, gradually moving downstream over the rearing period. As
2 they metamorphose, pacific lamprey move from fine substrate in lower
3 velocity waters into areas with silt covered gravels with greater water
4 velocity (Luzier et al. 2011). Juveniles move downstream, presumably
5 during high flow events in winter and spring. After reaching the ocean,
6 they spend approximately 3.5 years in saltwater (Beamish 1980). Adults
7 are parasitic, often found feeding on Chinook salmon, rockfish, and flatfish
8 (Luzier et al. 2011).

9 **River Lamprey** River lamprey (*Lampetra ayresi*) is designated as a State
10 species of special concern by DFG. River lamprey congregate upstream
11 from saltwater for 4 months as young adults, rapidly grow to 10 inches to
12 12 inches (25 centimeters to 31 centimeters), and enter the ocean in late
13 spring (Moyle 2002). After approximately 3 months in the ocean, river
14 lamprey return to freshwater to spawn in the fall (Moyle 2002). River
15 lamprey may hold in freshwater for up to 8 months until spawning from
16 April through June (Beamish 1980).

17 **Striped Bass** Striped bass are anadromous fish that have been an
18 important part of the sportfishing industry in the Delta. They were
19 introduced into the Bay-Delta between 1879 and 1882 (Moyle 2002).
20 Striped bass will not use fish ladders; therefore, their range in the
21 Sacramento River is limited to the reach of the river below RBDD. Striped
22 bass may move into the lower reaches of the major tributary rivers in the
23 Extended SPA year round, but probably most often between April and
24 June, when they spawn. The species tends to remain in deep, slow-moving
25 water, where it has access to prey without expending a great deal of energy.
26 Striped bass are a major predatory fish in the Delta, especially near human-
27 managed facilities, and are a source of mortality for delta smelt, juvenile
28 Chinook salmon and steelhead, and other fish species.

29 **Reservoir Fisheries** In most reservoirs, fish populations decline with the
30 aging of the reservoirs. For a variety of reasons, new reservoirs often
31 develop outstanding populations of fish that gradually decline as the
32 reservoirs mature. Loss of cover in the form of inundated vegetation is a
33 major contributing factor. Most Central Valley reservoirs are more than 25
34 years old, and these factors are likely reducing reservoir fish communities
35 from the population levels sustained shortly after these same reservoirs
36 were filled.

37 Mid-elevation reservoirs support a mixture of native fishes (species that
38 lived in the streams before dam construction) and introduced exotic
39 species. In many cases, the native species have become uncommon after an
40 initial 5–10 years of abundance. Over time, a variety of exotic species tends
41 to dominate the fish fauna in these reservoirs. The exact species

1 composition in each reservoir is related to the history of introductions, but
2 some species are almost universal in their occurrence: bluegill, largemouth
3 bass, carp, golden shiner, black crappie, brown bullhead, mosquitofish, and
4 rainbow trout (hatchery strains). A few native species, such as Sacramento
5 sucker and hitch, are permanently established in a number of Central
6 Valley reservoirs (Moyle 2002).

7 **3.5.2 Regulatory Setting**

8 The following text summarizes federal, State, and regional and local laws
9 and regulations pertinent to evaluation of the proposed program's impacts
10 on aquatic biological resources.

11 ***Federal***

12 **Endangered Species Act** The ESA protects and promotes recovery of
13 threatened and endangered species, many of which are aquatic and present
14 in the Extended SPA. Section 4 of the ESA outlines a process to list species
15 in danger of becoming extinct. Section 9 prohibits take of any threatened or
16 endangered species, including harm associated with habitat modifications.
17 Section 7 and Section 10 provide for exemptions on take prohibitions.
18 Under the ESA, the definition of "take" is to "harass, harm, pursue, hunt,
19 shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any
20 such conduct." USFWS and NMFS have also interpreted the definition of
21 "harm" to include significant habitat modification that could result in take.
22 If it is likely that implementing an action authorized, undertaken, or funded
23 by a federal agency would result in take of a federally listed species, a
24 federal interagency consultation, under Section 7 of the ESA, is required.
25 Additionally, some species may be in decline or at risk, but insufficient
26 information is available to indicate a need to list them as either threatened
27 or endangered; these species are listed as "species of concern." USFWS has
28 jurisdiction over terrestrial and nonanadromous fish species, and NMFS is
29 responsible for protecting anadromous fish and other marine species,
30 including marine mammals.

31 **Sustainable Fisheries Act (Essential Fish Habitat)** In response to
32 growing concern about the status of fisheries in the United States, Congress
33 passed the Sustainable Fisheries Act of 1996 (Public Law 104-297). This
34 law amended the Magnuson-Stevens Fishery Conservation and
35 Management Act (Public Law 94-265), the primary law governing marine
36 fisheries management in the federal waters of the United States. Under the
37 Sustainable Fisheries Act, consultation is required by NMFS on any
38 activity that might adversely affect Essential Fish Habitat (EFH). EFH
39 consists of those habitats that fish rely on throughout their life cycles. It
40 encompasses habitats necessary to allow sufficient production of

1 commercially valuable aquatic species to support a long-term sustainable
2 fishery and contribute to a healthy ecosystem.

3 **Fish and Wildlife Coordination Act** The Fish and Wildlife Coordination
4 Act, as amended in 1964, was enacted to protect fish and wildlife when
5 federal actions result in the control or modification of a natural stream or
6 body of water. The statute requires federal agencies to consider the effect
7 that water-related projects would have on fish and wildlife resources.
8 Federal agencies must consult and coordinate with State fish and game
9 agencies and USFWS to address ways to conserve wildlife resources by
10 preventing loss of and damage to these resources, and to further develop
11 and improve them. Adoption of the CVFPP is a State action and would not
12 trigger the need to comply with the Fish and Wildlife Coordination Act, but
13 related federal flood management actions must comply with this law.

14 **Executive Orders** The executive orders discussed below were issued to
15 provide direction to federal agencies regarding invasive species, floodplain
16 management, and protection of wetlands, and affect related federal flood
17 management actions.

- 18 • **Executive Order 13112: Invasive Species**—This executive order
19 directs federal agencies to prevent and control introductions of invasive
20 nonnative species in a cost-effective and environmentally sound
21 manner to minimize their economic, ecological, and human health
22 impacts. As directed by Executive Order 13112, a national invasive
23 species management plan guides federal actions to prevent, control, and
24 minimize invasive species and their impacts (NISC 2008). To support
25 implementation of this plan, the U.S. Army Corps of Engineers
26 (USACE) released the *U.S. Army Corps of Engineers Invasive Species*
27 *Policy* (USACE 2009). This policy calls on agencies to address the
28 effects of invasive species in impact analyses completed for civil works
29 projects.
- 30 • **Executive Order 11988: Floodplain Management**—This executive
31 order requires federal agencies to provide leadership and take action to
32 avoid development in the base (100-year) floodplain; reduce the
33 hazards and risk associated with floods; minimize the effect of floods
34 on human safety, health, and welfare; and restore and preserve the
35 natural and beneficial values of the base floodplain.
- 36 • **Executive Order 11990: Protection of Wetlands**—This executive
37 order directs federal agencies to provide leadership and act to minimize
38 the destruction, loss, or degradation of wetlands, and to preserve and
39 enhance the natural and beneficial values of wetlands in implementing
40 civil works.

1 **Central Valley Project Improvement Act** Implementing the CVPIA
2 changed management of the CVP by making protection of fish and wildlife
3 a project purpose, equal to water supply for agricultural and urban uses.
4 The CVPIA affects water exports from the Delta to San Luis Reservoir and
5 increases operational pressures on the reservoir to meet south-of-Delta
6 water demands. CVPIA Section 3406(b)(2) authorized and directed the
7 Secretary of the Interior to, among other actions, dedicate and manage 800
8 thousand acre-feet of CVP yield annually to meet the following goals:

- 9 • Implement the fish, wildlife, and habitat restoration purposes and
10 measures authorized in the CVPIA.
- 11 • Assist the State of California in its efforts to protect the waters of the
12 San Francisco Bay-Delta estuary.
- 13 • Help meet obligations legally imposed on the CVP under State or
14 federal law following the date of enactment of the CVPIA.

15 CVPIA Section 3406(d)(1) required the Secretary of the Interior to
16 immediately provide specific quantities of water (i.e., “Level 2” supplies)
17 to national wildlife refuges in the Central Valley. The CVPIA requires
18 delivery of Level 2 water in all year types except critically dry water years,
19 when Level 2 water can be reduced by 25 percent. Section 3406(d)(2) of
20 the CVPIA refers to “Level 4” refuge water supplies, the quantities
21 required for optimum habitat management of existing refuge lands. Level 4
22 water supplies amount to about 163 thousand acre-feet above Level 2 water
23 supplies. The availability of Level 4 refuge water supplies is influenced by
24 the availability of water for transfer from willing sellers.

25 CVPIA Section 3406(c)(1) mandated development of a reasonably prudent,
26 feasible comprehensive plan to be presented to Congress to address
27 concerns about fish, wildlife, and habitat on the San Joaquin River.
28 However, Public Law 111-11 declared that the 2006 settlement of a lawsuit
29 between several parties on restoration of the San Joaquin River, which led
30 to formation of the SJRRP, “satisfies and discharges all of the obligations
31 of the Secretary [of the Interior] contained in Section 3406(c)(1).”

32 **CALFED Bay-Delta Program** CALFED is a collaborative effort of 25
33 State and federal agencies focusing on restoring the ecological health of the
34 Bay-Delta, while ensuring improvements to water quality and reliability of
35 the water supply for all users of Bay-Delta water resources. CALFED
36 includes a range of balanced actions that are used in a comprehensive,
37 multiagency approach to managing Bay-Delta resources. The following are
38 CALFED’s objectives:

- 1 • Provide good water quality for all beneficial uses.
- 2 • Improve and increase aquatic and terrestrial habitats and improve
3 ecological functions in the Bay-Delta to support sustainable populations
4 of diverse and valuable plant and animal species.
- 5 • Reduce the mismatch between Bay-Delta water supplies and current
6 and projected beneficial uses dependent on the Bay-Delta system.
- 7 • Reduce the risk to land use and associated economic activities, water
8 supply, infrastructure, and the ecosystem from catastrophic breaching
9 of Delta levees.

10 The program objectives have been implemented among numerous
11 CALFED elements since the CALFED record of decision was certified in
12 2000.

13 **San Joaquin River Restoration Program** The SJRRP was formed in
14 response to the 2006 settlement of an 18-year-old lawsuit between the U.S.
15 Departments of the Interior and Commerce, the Natural Resources Defense
16 Council, and the Friant Water Users Authority. The Settlement establishes
17 two primary goals: (1) to restore and maintain fish populations in “good
18 condition” in the mainstem San Joaquin River below Friant Dam to the
19 confluence of the Merced River, including naturally reproducing and self-
20 sustaining populations of salmon and other fish; and (2) to reduce or avoid
21 adverse water supply impacts on all of the Friant Division long-term
22 contractors that may result from the Interim and Restoration flows provided
23 for in the Settlement.

24 The State has expressed strong support for this settlement and has pledged
25 cooperation and the State’s financial resources to help it succeed. The
26 settling parties and the California Resources Agency (now the California
27 Natural Resources Agency), DWR, and DFG entered into a memorandum
28 of understanding to allow the State to play a major, collaborative role in
29 planning, designing, funding, and implementing actions called for by the
30 Settlement Act to restore the San Joaquin River.

31 **National Wildlife Refuge Complex Comprehensive Conservation Plans**
32 USFWS’s San Luis National Wildlife Refuge (NWR) Complex consists of
33 San Luis NWR, Merced NWR, San Joaquin River NWR, and Grassland
34 Wildlife Management Area. These refuges comprise wetlands, grasslands,
35 riparian habitats, and agricultural fields, and many support fisheries
36 resources. The management goals and objectives for each refuge include
37 managing and providing habitat for endangered and sensitive species.
38 Those goals and objectives are set forth in 15-year comprehensive

1 conservation plans prepared by USFWS pursuant to the National Wildlife
2 Refuge System Improvement Act of October 1997.

3 **Clean Water Act** The objective of the Clean Water Act (CWA) is “to
4 restore and maintain the chemical, physical, and biological integrity of the
5 nation’s waters.” The CWA is the major federal legislation that governs
6 federal oversight of discharges into “jurisdictional waters” by federal, state,
7 local, and private activities. Jurisdictional waters are waters of the United
8 States, including wetlands. Waters of the United States include wetlands
9 and lakes, rivers, streams, and their tributaries. Wetlands are defined for
10 regulatory purposes in CWA Section 404 (described further below) as
11 “areas inundated or saturated by surface water or groundwater at a
12 frequency and duration sufficient to support and, under normal
13 circumstances do support, vegetation typically adapted for life in saturated
14 soil conditions.”

15 The CWA establishes the basic structure for regulating discharge of
16 pollutants into the waters of the United States and gives the U.S.
17 Environmental Protection Agency (EPA) the authority to implement
18 pollution control programs such as setting wastewater standards for
19 industries. In certain states such as California, EPA has delegated authority
20 to state agencies for most but not all CWA purposes.

21 *Section 303* Section 303 of the CWA requires states to adopt water quality
22 standards for all surface waters of the United States. Section 303(d) of the
23 CWA requires states and authorized Native American tribes to develop lists
24 of water quality–impaired segments of waterways. Each state’s or tribe’s
25 list identifies waters that do not meet the water quality standards necessary
26 to support the beneficial uses of a waterway, even after point sources
27 (identifiable localized sources of pollution) have installed the minimum
28 required pollution control technology. Only waters impaired by
29 “pollutants” (clean sediments, nutrients such as nitrogen and phosphorus,
30 pathogens, acids/bases, temperature, metals, cyanide, and synthetic organic
31 chemicals (EPA 2002)) are to be included on the list. Waters impaired by
32 other types of pollution (e.g., altered flow, channel modification) are not
33 included.

34 Section 303(d) of the CWA also requires states to maintain a list of
35 impaired water bodies so that a total maximum daily load (TMDL) can be
36 established. A TMDL is a plan to restore the beneficial uses of a stream or
37 to otherwise correct an impairment. It establishes the allowable pollutant
38 loadings or other quantifiable parameters (e.g., pH, temperature) for a
39 water body, thereby providing the basis for establishing water quality–
40 based controls. The calculation used to establish TMDLs for a water body
41 must include a margin of safety to ensure that the water body can be used

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 for the purposes of State designation. The calculation also must account for
2 seasonal variation in water quality (EPA 2002). The Central Valley
3 Regional Water Quality Control Board (Central Valley RWQCB) develops
4 TMDLs for the Sacramento and San Joaquin rivers, and for many
5 tributaries associated with these rivers (see the discussion of the Porter-
6 Cologne Water Quality Control Act (Porter-Cologne Act) below).

7 *Sections 401, 402, and 404* Section 401 of the CWA regulates the water
8 quality aspects of a proposed activity, which also affect fisheries and other
9 aquatic resources. Section 401 certification is the responsibility of the
10 SWRCB and the appropriate RWQCB (in this case, the Central Valley
11 RWQCB), which certifies that the activity is consistent with State-issued
12 water quality control plans, called basin plans. Section 401 also requires
13 federal agencies to obtain certification from the State or Native American
14 tribes before issuing permits that would result in increased pollutant loads
15 to a water body. The certification is issued only if such increased loads
16 would not cause or contribute to exceedences of water quality standards.

17 Section 402 created the National Pollutant Discharge Elimination System
18 permitting program. This program covers discharges of point sources of
19 pollution, including stormwater discharges, into a surface water body.

20 Section 404 of the CWA regulates the discharge of dredged and fill
21 material in jurisdictional waters. Specifically, a permit must be obtained
22 from USACE under Section 404 for the discharge of dredged or fill
23 material into waters of the United States, including wetlands (defined
24 above). Under the 404 permit process, certain activities, such as
25 maintenance, can be covered by nationwide permits; other activities, such
26 as emergency response, are covered by regional permits. Activities not
27 covered under either nationwide or regional permits need to be addressed
28 through an individual permit process. Water quality certification under
29 Section 401 of the CWA is required for all projects receiving Section 404
30 permits.

31 **Rivers and Harbors Act, Sections 10 and 408** Section 10 of the Rivers
32 and Harbors Act of 1899 (33 U.S. Code (USC) 401 et seq.) requires project
33 proponents to obtain authorization from USACE before constructing any
34 structure over, in, or under navigable waters of the United States. Under
35 Section 14 of the Rivers and Harbors Act (33 USC 408)—most often
36 referred to as Section 408—the Secretary of the Army is authorized to
37 permit alterations/modifications to existing USACE project levees in
38 certain circumstances. The Secretary of the Army has delegated this
39 approval authority to the USACE Chief of Engineers. The authority to
40 approve relatively minor, low-impact alterations/modifications related to
41 operational and maintenance-related responsibilities of the nonfederal

1 entities responsible for the flood control system (e.g., the Central Valley
2 Flood Protection Board (Board), local reclamation districts) has been
3 further delegated to the applicable USACE District Engineer, in accordance
4 with 33 CFR 208.10.

5 Placement of structures such as pump houses, stairs, pipes, bike trails,
6 sidewalks, fences, driveways, power poles, and instrumentation can be
7 approved by a District Engineer, provided that these
8 alterations/modifications do not adversely affect the functioning of the
9 project and flood-fighting activities. The types of alterations/modifications
10 that require approval by the Chief of Engineers under 33 USC 408 include
11 degradations, raisings, and realignments and other alteration/modifications
12 to the flood protection system not discussed above. In administering
13 Sections 10 and 408, USACE must consider the environmental effects of
14 actions regulated under these statutes, especially with respect to aquatic
15 resources and fisheries.

16 For activities in the SPFC, the Board acts as the nonfederal sponsor. In this
17 capacity, the Board coordinates reviews and submits project requests,
18 project designs, and technical engineering documents to USACE for
19 consideration under 33 USC 408 and 33 USC 208.10.

20 **Federal Power Act, Section 18** Section 18 of the Federal Power Act
21 authorizes the U.S. Departments of the Interior and Commerce to require
22 fishways in new licenses granted by the Federal Energy Regulatory
23 Commission (discussed further in Section 3.9, “Energy”) when it can be
24 demonstrated that fish populations would benefit from the provision of fish
25 passage.

26 **State**

27 **California Water Code** The California Water Code authorizes the
28 SWRCB to allocate surface water rights, and to permit diversion and use of
29 water throughout California. The SWRCB considers effects on fisheries as
30 part of its permitting process. Division 7 of the California Water Code,
31 known as the Porter-Cologne Act, regulates activities that affect water
32 quality (see the separate discussion of the Porter-Cologne Act below).
33 Division 5 of the Water Code specifies roles and responsibilities for flood
34 control for the State and for numerous local agencies throughout California.

35 **California Endangered Species Act** As part of the CESA and Section
36 2081 of the California Fish and Game Code, a permit from DFG is required
37 for projects that could result in the taking of a species that is State-listed as
38 threatened or endangered. Under the CESA, “take” is defined as an activity
39 that would directly or indirectly kill an individual of a species; however,
40 the CESA definition does not include “harm” or “harass,” as the federal

1 ESA definition does. As a result, the threshold for take is higher under the
2 CESA than under the federal ESA.

3 **California Fish and Game Code** The sections of the California Fish and
4 Game Code listed below provide environmental protections and could
5 apply to specific CVFPP projects as they are defined and proposed.

- 6 • **Section 1602, Streambed Alteration**—Diversions, obstructions, or
7 changes to the natural flow or bed, channel, or bank of any river,
8 stream, or lake in California that supports wildlife resources are subject
9 to regulation by DFG.
- 10 • **Fully Protected Species under the California Fish and Game**
11 **Code**—Protection of fully protected species is described in four
12 sections of the California Fish and Game Code that list 37 fully
13 protected species (Sections 3511, 4700, 5050, and 5515), many of
14 which are present in the Extended SPA. These statutes prohibit take or
15 possession at any time of fully protected species.
- 16 • **Section 5937**—Under most conditions, sufficient volumes of water are
17 required to pass through a fishway at all times. In the absence of a
18 fishway, sufficient water must be allowed to pass over, around, or
19 through a dam to keep in good condition any fish that may be planted
20 or exist below the dam.

21 **California Code of Regulations, Title 23** Under Title 23 of the
22 California Code of Regulations, the Board cooperates with local, State, and
23 federal agencies in establishing, planning, constructing, operating, and
24 maintaining flood control works in the Central Valley. The Board is
25 required to enforce appropriate standards for constructing, maintaining, and
26 protecting adopted flood control plans that will best protect the public from
27 floods along the Sacramento and San Joaquin rivers and their tributaries.
28 The Board issues encroachment permits to maintain the integrity and safety
29 of flood control project levees and floodways.

30 **Porter-Cologne Water Quality Control Act** The Porter-Cologne Act is
31 California's statutory authority for the protection of water quality. Under
32 the act, the State must adopt water quality policies, plans, and objectives
33 protecting the State's waters for the use and enjoyment of the people. The
34 act sets forth the obligations of the SWRCB and RWQCBs to adopt and
35 periodically update their basin plans. A basin plan identifies the designated
36 beneficial uses for specific surface water and groundwater resources,
37 applicable water quality objectives necessary to support the beneficial uses,
38 and implementation programs that are established by the RWQCBs to
39 maintain and protect water quality from degradation. The Porter-Cologne

1 Act also requires waste dischargers to notify the RWQCBs of their
2 activities by filing reports of waste discharge. In addition, the act authorizes
3 the SWRCB and RWQCBs to issue and enforce waste discharge
4 requirements, National Pollutant Discharge Elimination System permits,
5 Section 401 water quality certifications, or other approvals. The RWQCBs
6 also have the authority to issue waivers to reports of waste discharge/waste
7 discharge requirements for broad categories of “low threat” discharge
8 activities that have minimal potential for adverse effects on water quality,
9 when implemented according to prescribed terms and conditions.

10 **State Lands Commission** The State Lands Commission has exclusive
11 jurisdiction over all ungranted tidelands and submerged lands owned by the
12 State, and the beds of navigable rivers, sloughs, and lakes. A project cannot
13 use these State lands unless a lease is first obtained from the State Lands
14 Commission.

15 ***Regional and Local***

16 **Sacramento River Watershed Program** The Sacramento River
17 Watershed Program (SRWP), founded in 1996, brings together dozens of
18 groups and thousands of people who are concerned about the health of the
19 Sacramento River and its watershed. As one of the largest watersheds in the
20 United States, the Sacramento River watershed serves as an important
21 source of drinking water and recreation, as well as a vital economic artery
22 for commerce and agriculture. Therefore, preserving and maintaining the
23 water quality of the Sacramento River watershed is crucial. The program is
24 overseen by a 21-member board of trustees and functions through several
25 committees and work groups.

26 The program provides a network for building a basinwide context to
27 improve watershed health. It operates through consensus-based
28 collaborative partnerships, coordination of research and monitoring, and
29 mutual education among the stakeholders of the Sacramento River
30 watershed. The SRWP works to support and preserve the integrity of local
31 efforts. The program strives to resolve watershed issues with local
32 participation and a watershed-wide perspective. The SRWP also helps
33 disseminate information about the watershed and conducts monitoring
34 activities to continually assess water quality and other indicators of
35 watershed health.

36 **Lower Yuba River Accord** The Lower Yuba River Accord (Yuba
37 Accord) enables Yuba County Water Agency to successfully operate the
38 Yuba River Development Project (Federal Energy Regulatory Commission
39 Project No. 2246, 362 megawatts) for hydropower, irrigation, flood control,
40 recreation, and fisheries benefits. As a settlement agreement, the Yuba
41 Accord is the final product of negotiations among stakeholders, which

1 include local irrigation districts, State and federal resource agencies, and
2 conservation groups. The State approved the agreement in 2008, and the
3 project is now fully operational. The Yuba Accord is unprecedented in that
4 it combines increased instream flows for wild, native salmon and steelhead
5 with increased supplemental water supplies for California cities and farms,
6 while preserving the project's capacity to generate clean, renewable
7 hydropower. The Yuba Accord also reaffirms the water rights of Yuba
8 County Water Agency and its member irrigation districts.

9 **Lower American River Corridor Management Plan** The *Lower*
10 *American River Corridor Management Plan* serves to promote a
11 cooperative approach to managing and enhancing the lower American
12 River within the framework of the *American River Parkway Plan 2008*
13 (Sacramento County 2008). The goals of the river corridor management
14 plan are to protect and enhance fisheries and instream habitat, protect and
15 enhance vegetation and wildlife habitat, improve the reliability of the
16 existing flood control system, and enhance the lower American River's
17 wild and scenic recreation values (Lower American River Task Force
18 2002).

19 **Bay Delta Conservation Plan** The *Bay Delta Conservation Plan* (BDCP)
20 is being developed in compliance with the federal ESA and the California
21 Natural Communities Conservation Planning Act. When complete, the
22 BDCP will provide the basis for the issuance of endangered species permits
23 for the operation of the State and federal water projects. Once approved,
24 the plan would be implemented over the next 50 years. The heart of the
25 BDCP is a long-term conservation strategy that sets forth actions needed
26 for a healthy Delta and alternative water conveyance. The BDCP is being
27 prepared through collaboration among local, State, and federal water
28 agencies; State and federal fish and wildlife agencies; environmental
29 organizations; and other interested parties. These organizations have
30 formed the BDCP Steering Committee, with the goal of identifying water
31 flow and habitat restoration actions to recover endangered and sensitive
32 species and their habitats in California's Delta. The lead agencies for the
33 BDCP environmental impact report/environmental impact statement are
34 DWR, Reclamation, USFWS, and NMFS; these agencies are acting in
35 cooperation with DFG, EPA, and USACE.

36 **San Joaquin River Parkway Master Plan** The *San Joaquin River*
37 *Parkway Master Plan* is a regional resource management plan for the San
38 Joaquin River area between Friant Dam and State Route 99. The San
39 Joaquin River Conservancy, a regionally governed agency created by the
40 State, is charged with implementing this master plan. The plan's main
41 tenets are the protection of natural resources, public education, and the

1 promotion of low-impact recreational use of the river corridor (SJRC
2 2000).

3 **Lower Feather River Corridor Management Plan** The *Lower Feather*
4 *River Corridor Management Plan* is being developed by DWR to establish
5 a vision for future management, restoration, and maintenance of flood
6 control facilities, conveyance channels, and floodplain and related habitat
7 on the Feather River from the Sutter Bypass to the Yuba River confluence
8 (approximately 20 miles). The plan will implement the new collaborative
9 approach for planning, designing, and implementing projects within and
10 adjacent to flood control features that DWR is responsible for maintaining
11 and repairing.

12 **County and City Policies and Ordinances** Numerous counties and cities
13 throughout the study area have established a multitude of policies and
14 ordinances that address local protection of fisheries, sensitive species, and
15 aquatic resources; many of them are applicable to the CVFPP. Should a
16 place-based project be defined and pursued as part of the proposed
17 program, and should the CEQA lead agency be subject to the authority of
18 local jurisdictions, the applicable county and city policies and ordinances
19 would be addressed in a project-level CEQA document as necessary.

20 **3.5.3 Analysis Methodology and Thresholds of** 21 **Significance**

22 This section provides a program-level evaluation of the direct and indirect
23 effects on aquatic resources of implementing management actions included
24 in the proposed program. These proposed management actions are
25 expressed as NTMAs and LTMAAs. The methods used to assess how
26 different categories of NTMAs and LTMAAs could affect aquatic resources
27 are summarized in “Analysis Methodology”; thresholds for evaluating the
28 significance of potential impacts are listed in “Thresholds of Significance.”
29 Potential effects related to each significance threshold are discussed in
30 Section 3.5.4, “Environmental Impacts and Mitigation Measures for
31 NTMAAs,” and Section 3.5.5, “Environmental Impacts, Mitigation
32 Measures, and Mitigation Strategies for LTMAAs.”

33 **Analysis Methodology**

34 Impact evaluations were based on a review of the management actions
35 proposed under the CVFPP, expressed as NTMAAs and LTMAAs in this
36 PEIR, to determine whether these actions could potentially result in
37 impacts on aquatic resources. NTMAAs and LTMAAs are described in more
38 detail in Section 2.4, “Proposed Management Activities.” The overall
39 approach to analyzing the impacts of NTMAAs and LTMAAs and providing
40 mitigation is summarized below and described in detail in Section 3.1,

1 “Approach to Environmental Analysis.” NTMAs can consist of any of the
2 following types of activities:

- 3 • Improvement, remediation, repair, reconstruction, and operation and
4 maintenance of existing facilities
- 5 • Construction, operation, and maintenance of small setback levees
- 6 • Purchase of easements and/or other interests in land
- 7 • Operational criteria changes to existing reservoirs that stay within
8 existing storage allocations
- 9 • Implementation of the vegetation management strategy (VMS) included
10 in the CVFPP
- 11 • Initiation of conservation elements included in the proposed program
- 12 • Implementation of various changes to DWR and Statewide policies that
13 could result in alteration of the physical environment

14 All other types of CVFPP activities fall within the LTMA category.
15 NTMAs are evaluated using a typical “impact/mitigation” approach. Where
16 impact descriptions and mitigation measures identified for NTMAs also
17 apply to LTMAs, they are also attributed to LTMAs, with modifications or
18 expansions as needed. However, because many LTMAs are more general
19 and conceptual, additional impacts are described in a broader narrative
20 format. Impacts of LTMAs that are addressed in this narrative format are
21 those considered too speculative for detailed evaluation, consistent with
22 Section 15145 of the CEQA Guidelines. Following the narrative
23 description of these additional impacts is a list of suggested mitigation
24 strategies that could be employed, indicating the character and scope of
25 mitigation actions that might be implemented if a future project-specific
26 CEQA analysis were to find these impacts to be significant.

27 ***Thresholds of Significance***

28 The following applicable thresholds of significance have been used to
29 determine whether implementing the proposed program would result in a
30 significant impact. These thresholds of significance are based on Appendix
31 G of the CEQA Guidelines, as amended. An impact on aquatic resources is
32 considered significant if implementation of the proposed program would do
33 any of the following when compared against existing conditions:

- 34 • Have a substantial adverse effect, either directly or through habitat
35 modifications, on any fish species identified as a candidate, sensitive, or

- 1 special-status species in local or regional plans, policies, or regulations,
2 or by DFG, NMFS, or USFWS
- 3 • Interfere substantially with the movement of any native resident or
4 migratory fish species or impede the use of native fish nursery sites
 - 5 • Have a substantial adverse effect on riparian vegetation that functions
6 as shaded riverine aquatic habitat
 - 7 • Substantially reduce the habitat of a fish species; cause a population to
8 drop below self-sustaining levels; or threaten to eliminate a fish
9 community
 - 10 • Reduce the number or restrict the range of a rare or endangered aquatic
11 species
 - 12 • Substantially reduce habitat designated as critical habitat or Essential
13 Fish Habitat (EFH)
 - 14 • Substantially conflict with the provisions of an adopted habitat
15 conservation plan (HCP), natural community conservation plan, or
16 other approved local, regional, or State HCP

17 The substantial effects referred to in these criteria may occur through
18 various mechanisms, such as modification of riparian vegetation and
19 riverine habitat; long-term modification of channels or reservoirs (e.g.,
20 storage); and long-term reduction in the acreage of federally protected
21 waters of the United States capable of supporting aquatic species. See
22 Section 3.6, “Biological Resources—Terrestrial,” for additional discussions
23 of riparian and wetland impacts.

24 **3.5.4 Environmental Impacts and Mitigation Measures** 25 **for NTMAs**

26 This section describes the physical effects of NTMAs on aquatic biological
27 resources. For each impact discussion, the environmental effect is
28 determined to be either less than significant, significant, potentially
29 significant, or beneficial compared to existing conditions and relative to the
30 thresholds of significance described above. These significance categories
31 are described in more detail in Section 3.1, “Approach to Environmental
32 Analysis.” Feasible mitigation measures are identified to address any
33 significant or potentially significant impacts. Actual implementation,
34 monitoring, and reporting of the PEIR mitigation measures would be the
35 responsibility of the project proponent for each site-specific project. For
36 those projects not undertaken by, or otherwise subject to the jurisdiction of,
37 DWR or the Board, the project proponent generally can and should

1 implement all applicable and appropriate mitigation measures. The project
2 proponent is the entity with primary responsibility for implementing
3 specific future projects and may include DWR; the Board; reclamation
4 districts; local flood control agencies; and other federal, State, or local
5 agencies. Because various agencies may ultimately be responsible for
6 implementing (or ensuring implementation of) mitigation measures
7 identified in this PEIR, the text describing mitigation measures below does
8 not refer directly to DWR but instead refers to the “project proponent.”
9 This term is used to represent all potential future entities responsible for
10 implementing, or ensuring implementation of, mitigation measures.

11 **Impact BIO-A-1 (NTMA): *Potential Effects on Special-Status Fish,***
12 ***Fish Movement, Nursery Ground Usage, Riparian Habitat, Designated***
13 ***Critical Habitat, and Essential Fish Habitat Caused by Siltation and***
14 ***Degradation of Water Quality during Construction or Operations and***
15 ***Maintenance Activities***

16 Any erosion resulting from construction or operations and maintenance
17 activities (particularly with respect to levees) required for NTMAs could
18 temporarily increase turbidity and sedimentation downstream from the
19 construction sites if soils were to be transported during in-water work or in
20 stormwater runoff. Sedimentation and increased turbidity or other
21 contamination could degrade water quality and adversely affect fish habitat
22 (riparian habitat, critical habitat, and EFH), fish movement, and fish
23 populations (including special-status fish species).

24 Fish population levels and survival have been linked to levels of turbidity
25 and siltation in a watershed. Prolonged exposure to high levels of
26 suspended sediment can create a loss of visual capability, leading to a
27 reduction in feeding and growth rates; a thickening of the gill epithelium,
28 potentially causing the loss of respiratory function; clogging and abrasion
29 of gill filaments; and increases in stress levels, reducing the tolerance of
30 fish to disease and toxicants (Waters 1995).

31 High levels of suspended sediment also cause the movement and
32 redistribution of fish populations, and can affect physical habitat. Once
33 suspended sediment is deposited, it can reduce water depths in pools,
34 decreasing the water’s physical carrying capacity for juvenile and adult fish
35 (Waters 1995) and thereby decreasing the quality of critical habitat or EFH.
36 Increased sediment loading can degrade food-producing habitat
37 downstream from the project area as well. Sediment loading can interfere
38 with photosynthesis of aquatic flora and displace aquatic fauna. Many fish
39 are sight feeders, and turbid waters reduce the efficiency of these fish in
40 locating and feeding on prey. Some fish, particularly juveniles, can become
41 disoriented and leave areas where their main food sources are located,

1 ultimately reducing their growth rates. Additionally, benthic
2 macroinvertebrates, a main food source for numerous fishes, can be found
3 in much lower densities in highly turbid areas.

4 Avoidance behavior is the most common result of increases in turbidity and
5 sedimentation. Fish will not occupy areas that are not suitable for survival
6 unless they have no other option. Therefore, habitat can become limited in
7 systems where high turbidity precludes a species from occupying habitat
8 required for specific life stages.

9 The potential also exists for contaminants such as concrete, fuels, oils, and
10 other petroleum products used in construction activities to be introduced in
11 the water system, either directly or through surface runoff. Contaminants
12 may be toxic to fish and benthic macroinvertebrates or may change oxygen
13 diffusion rates and cause acute and chronic toxicity to aquatic organisms,
14 thereby reducing growth and survival.

15 Project proponents or others implementing construction activities must file
16 a notice of intent with the Central Valley RWQCB to discharge stormwater
17 associated with construction activity. As part of the final design and
18 construction specifications for NTMAs, project proponents or others
19 implementing construction activities would be required to implement
20 standard best management practices (BMPs) related to erosion, siltation,
21 and “good housekeeping.” Before implementing NTMAs, construction
22 contractors would be required to prepare and implement storm water
23 pollution prevention plans (SWPPPs) and comply with the conditions of
24 the National Pollutant Discharge Elimination System general stormwater
25 permit for construction activity (Order No. 2009-0009-DWQ). The
26 SWPPPs would describe the construction activities to be conducted, BMPs
27 to be implemented to prevent discharges of contaminated stormwater into
28 waterways, and inspection and monitoring activities to be conducted.

29 The SWPPP for each NTMA would include pollution prevention measures;
30 a demonstration of compliance with all applicable standards of the Central
31 Valley RWQCB and other relevant water quality, erosion, and sediment
32 control standards; checklists for maintenance inspections; detailed
33 construction timelines; and a BMP monitoring and maintenance schedule.
34 BMPs would include requirements to conduct all work according to site-
35 specific construction plans; install silt fences near riparian areas and
36 existing drainages; reseed cleared areas with native vegetation and stabilize
37 disturbed soils before the onset of the winter rainfall season; conduct
38 maintenance on a regular basis; and immediately repair and replace BMPs
39 that have failed.

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 The SWPPP also would specify appropriate handling, storage, and spill
2 response practices for hazardous materials to reduce the possibility of
3 adverse impacts from use, accidental spills, or releases of contaminants.
4 BMPs would be applied to meet the “maximum extent practicable” and
5 “best conventional technology/best available technology” requirements and
6 to address compliance with water quality standards. A monitoring program
7 would be implemented during and after construction to ensure that the
8 NTMA would comply with all applicable standards and that the BMPs
9 would be effective.

10 Where in-water work might be necessary, these activities would also
11 require authorization from USACE under Section 404 of the CWA and
12 from the appropriate RWQCB under Section 401 of the CWA. Permit
13 conditions from these agencies would include further actions to monitor
14 and protect water quality, such as using construction methods that
15 minimize soil disturbance and contain sediment via silt curtains or other
16 means.

17 Project proponents and others implementing construction activities must
18 develop and implement a SWPPP to avoid increased sedimentation and
19 turbidity and/or release of contaminants that could degrade aquatic habitats
20 and adversely affect aquatic species. They could be required to implement
21 additional measures if in-water work is required. Therefore, this impact
22 would be **less than significant**. No mitigation is required.

23 **Impact BIO-A-2 (NTMA): *Effects on Special-Status Fish, Fish***
24 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
25 ***Critical Habitat, and Essential Fish Habitat Caused by Loss of Overhead***
26 ***Cover and Instream Woody Material as Part of the Vegetation***
27 ***Management Strategy***

28 Implementing the VMS (as described in Section 2.4.3, “Other Near-Term
29 Management Activities,” in Chapter 2.0, “Program Description”) would
30 result in a gradual reduction of existing riparian habitats in some locations
31 on and along existing levees, as dead or diseased trees are removed and not
32 replaced by either natural recruitment or planting. Trees and other woody
33 vegetation would be removed over an extended period—and eventually
34 eliminated entirely—from the designated vegetation management zone, an
35 area typically extending 15 feet beyond the landside levee toe to 20 feet
36 below the waterside levee crown. Immature trees and woody vegetation
37 would be removed, existing mature trees either would be lost eventually to
38 natural mortality or would be removed if they posed an unacceptable threat,
39 and new trees and woody vegetation would not be reestablished. However,
40 vegetation would generally be retained on the water side of levees more
41 than 20 feet below the levee crown.

1 Specifically, under the VMS, immature trees and woody vegetation in the
2 vegetation management zone that measure less than 4 inches in diameter at
3 breast height (dbh) would be removed in an authorized manner as part of
4 levee maintenance. Larger trees and woody vegetation greater than 4 inches
5 dbh would be subject to a long-term life-cycle management (LCM) plan to
6 be implemented by levee maintenance agencies. These larger trees would
7 be allowed to live out their normal life cycles if they do not pose an
8 unacceptable threat, but would not be replaced in the vegetation
9 management zone after their death or removal. (The LCM plan allows the
10 immediate removal of trees that pose an unacceptable threat.) Removal of
11 woody vegetation in both size categories would be conducted in
12 consultation with the appropriate resource agencies.

13 Over time, a net loss in the extent and quality of riparian habitat would
14 occur in the vegetation management zone on existing levees as the lost
15 vegetation is not replaced. Vegetation less than 4 inches dbh would be
16 removed relatively quickly after plan adoption. Larger riparian vegetation
17 (e.g., mature cottonwoods and black willows) is expected to gradually
18 decline, and the vegetation management zone would ultimately consist
19 almost exclusively of smaller, nonwoody vegetation. Overhanging
20 vegetation, most often from large trees, provide stream shade, which is a
21 component of shaded riverine aquatic habitat.

22 The effects of vegetation removal under the VMS would vary substantially
23 depending on the existing conditions along a particular levee segment:

- 24 • In locations where little to no woody vegetation grows in the vegetation
25 management zone, and existing levee maintenance practices prevent
26 this vegetation from establishing, the VMS would result in little change
27 from existing conditions.
- 28 • If the ordinary water level approaches the waterside edge of the
29 vegetation management zone, and the only woody riparian vegetation
30 on the waterside of the levee is a thin strip in the management zone (20
31 feet or less below the crown), much of the woody riparian vegetation on
32 this side of the levee could be removed over time.
- 33 • If woody riparian vegetation grows on the levee's waterside both in and
34 below the vegetation management zone, riparian vegetation would be
35 lost in the management zone but retained below it. As a result, the strip
36 of waterside riparian habitat would be thinner than under existing
37 conditions.
- 38 • In situations where woody riparian vegetation grows on both sides of a
39 levee, and with some vegetation in the vegetation management zone,

1 the current nonriparian corridor between the landside and waterside
2 riparian vegetation (likely a levee crown patrol road and portions of the
3 levee slope) would become wider as vegetation in the management
4 zone on both sides of the levee moves toward an increased amount of
5 smaller and nonwoody vegetation.

6 Numerous other vegetation removal scenarios could be described here.
7 However, the key point is that as the VMS is implemented, adverse effects
8 on riparian vegetation and associated aquatic resources could range from
9 minimal to substantial, depending on factors such as location, amount, and
10 quality of vegetation affected; its proximity to water; and the continuity
11 with other riparian vegetation. Where adverse effects are found, they would
12 result primarily from one of three scenarios:

13 (1) Thin strips of riparian vegetation that grow entirely within the
14 vegetation management zone would be substantially or entirely
15 removed.

16 (2) Riparian vegetation grows both inside and outside of the vegetation
17 management zone, and habitat in the management zone ultimately
18 would be removed. As a result, thinner corridors of riparian habitat
19 would remain outside of the management zone.

20 (3) Woody riparian habitat exists on both sides of the levee, separated by a
21 nonriparian zone along the levee (likely, at a minimum, along a crown
22 patrol road). If some riparian habitat occurs within the vegetation
23 management zone, this habitat would be removed over time, causing
24 the nonriparian zone between the landside and waterside habitat to
25 become wider. However, this mechanism would be very unlikely to
26 affect aquatic resources, and potential adverse effects would typically
27 be limited to terrestrial biological resources. (See Section 3.6,
28 “Biological Resources—Terrestrial.”)

29 However, a component of both the VMS and the CVFPP Conservation
30 Framework is also the enhancement of existing riparian habitats and
31 restoration and creation of riparian habitat at various locations. Riparian
32 forest corridors would be established, as appropriate, in areas outside the
33 vegetation management zone along both the waterside and landside of
34 existing levees. The greatest opportunities to increase the extent of riparian
35 vegetation would be on the landside because of space limitations often
36 found between levees and the water bodies they are designed to contain. It
37 is most likely that restoration and creation of riparian forest corridors
38 would be in proximity to levees in rural areas where undeveloped land is
39 available and human disturbance would be minimized.

1 The VMS would also inform the design of new setback levees by
2 recommending an expanded floodway that would accommodate both
3 vegetation and water conveyance. Under this approach, woody vegetation
4 may be permitted on the waterside slopes and berms of new levees where a
5 specifically designed waterside planting berm is incorporated into the levee
6 design. In some cases, woody vegetation provides environmental and
7 engineering benefits to levee integrity (e.g., erosion protection, soil
8 reinforcement, sediment recruitment). In these cases, the vegetation could
9 remain on existing levees that are repaired or improved, particularly where
10 the levee prism is widened or a root or seepage barrier is installed. With
11 these efforts, existing riparian habitat could be retained or expanded along
12 levees at some locations.

13 The combined elements of the VMS would result in the removal of riparian
14 vegetation in some areas and the enhancement, restoration, or creation of
15 riparian vegetation in other areas. The final result would be a gradual
16 change in the location of riparian vegetation, with habitat lost in some areas
17 but gained in other areas. There is the potential that ultimately a net gain in
18 riparian vegetation could result. The recovery and restoration of native
19 habitats is a supporting goal of the CVFPP, and increasing and improving
20 the quantity, diversity, quality, and connectivity of riverine habitats
21 (including riparian habitat) is a goal of the Conservation Framework.
22 However, there is currently insufficient detail in these plans to ensure that,
23 in all time periods and in all areas, there would be a balance between
24 habitat losses and gains, resulting in no net overall loss in the extent and
25 quality of riparian vegetation in the program area relative to existing
26 conditions.

27 With the CVFPP Conservation Framework, planting riparian vegetation
28 below the vegetation management zone could enhance existing riparian
29 habitats and result in restoration or creation of additional riparian habitat at
30 various locations. A portion of the affected riparian habitat—both the gains
31 (below the vegetation management zone) and the losses (in and below the
32 vegetation management zone, if a matter of public safety)—may qualify as
33 shaded riverine aquatic habitat. This is an important habitat component for
34 aquatic species, including special-status fish species. Shaded riverine
35 aquatic habitat is also considered part of the critical habitat and EFH
36 particularly for salmonid species.

37 The effect of implementing the VMS (i.e., LCM) would be gradual for
38 woody vegetation greater than 4 inches dbh. Therefore, the rate at which
39 these habitat components would be enhanced, restored, and created under
40 the CVFPP Conservation Framework could match or exceed the rate of
41 potential habitat loss associated with the VMS. Ultimately, habitat
42 improvements resulting from implementation of the Conservation

1 Framework would likely exceed losses resulting from implementation of
2 the VMS on a net basis. The final outcome would be a gradual change in
3 the locations of riparian/shaded riverine aquatic vegetation as habitat is lost
4 in some areas but gained in other areas.

5 It cannot be assured that habitat gains generated by the CVFPP
6 Conservation Framework would always exceed losses at a specific
7 location. If vegetation removal were required in a general area that
8 currently has a high volume of riparian vegetation, the removal and offsite
9 mitigation would have less of an effect on the overall system because
10 changes in overall conditions in the area would be small. However, if
11 vegetation were removed in an area where minimal riparian vegetation is
12 available, this removal—even with offsite mitigation—would have a
13 greater effect on the fisheries. The effect would be greater because it is
14 more likely that connectivity between patches of riparian habitat could be
15 limited and long stretches of river shoreline would have little to no riparian
16 vegetation. Although clearly not every levee segment in the SPFC contains
17 riparian vegetation that functions as SRA habitat, it is reasonable to assume
18 that there would be some areas where SRA currently exists along relatively
19 long river reaches where this habitat would be removed. Therefore,
20 implementation of the VMS could have, at least in some areas, a substantial
21 adverse effect on riparian vegetation that functions as SRA habitat.

22 Because overhead cover and IWM (and thus shaded riverine aquatic
23 habitat) would be lost as a result of implementation of the VMS along the
24 banks and levees, this impact would be **potentially significant**.

25 **Mitigation Measure BIO-A-2a (NTMA): *Secure Applicable State and/or***
26 ***Federal Permits and Implement Permit Requirements***

27 Not all measures listed below may be applicable to each management
28 action. Rather, these measures serve as an overlying mitigation framework
29 to be used for specific management actions. The applicability of measures
30 listed below would vary based on the lead agency, location, timing, and
31 nature of each management action.

32 The project proponent will ensure that the following measures are
33 implemented to reduce the effects of repairing, reconstructing, and
34 improving levees on trees within stream zones, shaded riverine aquatic
35 habitat, IWM, listed fish species, and designated critical habitat:

- 36 • A Section 1602 streambed alteration agreement will be obtained from
37 DFG before any trees are removed from a stream zone that is under
38 DFG jurisdiction. The project proponent will comply with all terms and

1 conditions of the streambed alteration agreement, including measures to
2 protect habitat or to restore, replace, or rehabilitate any habitat.

- 3 • The project proponent will consult or coordinate with USFWS and
4 NMFS as required under the federal ESA, and with DFG as required
5 under the CESA, regarding potential impacts on listed fish species,
6 including the loss of habitat. The project proponent will implement any
7 additional measures developed through the ESA and CESA
8 consultation processes, including the conditions of Section 7 biological
9 opinions, Section 10 HCPs, and Section 2081 permits.

10 Where an existing approved HCP, NCCP, or similar plan covers an NTMA
11 and provides for compliance with applicable State or federal regulations,
12 the project proponent may participate in and comply with the terms of such
13 a plan to achieve the permit compliance measures listed above. Any
14 mitigation plantings in the floodway will not be permitted if they would
15 result in substantial increases in flood stage elevations.

16 **Mitigation Measure BIO-A-2b (NTMA): *Ensure Full Compensation for***
17 ***Losses of Riparian Habitat Functions and Values Caused by***
18 ***Implementing the Vegetation Management Strategy Along Levees***

19 DWR will coordinate with the Board and levee maintenance agencies
20 tasked with implementing the VMS to develop and implement a plan to
21 record data on riparian vegetation lost or removed due to implementation of
22 the VMS, and to ensure adequate compensation for losses of riparian
23 habitat functions and values. Although this mitigation measure is written as
24 if a single plan is prepared, multiple plans addressing individual regions,
25 watersheds, river corridors, or other geographic subdivisions are also
26 acceptable.

27 The plan will be completed and suitable for implementation before the start
28 of riparian habitat removal under the VMS. The plan will include
29 mechanisms to, at a minimum, record and track the acreage, type, and
30 location of riparian habitat to be removed through implementation of the
31 VMS or lost over time through LCM.

32 The plan will also address compensation for the loss and degradation of
33 riparian habitat through the enhancement, restoration, or creation of
34 riparian habitat in other locations. Assessment of the value of lost or
35 degraded habitat and of compensation habitat will take into account issues
36 such as the differing functions of waterside and landside riparian habitat,
37 continuity and connectivity of habitat, types of riparian habitat removed vs.
38 type of compensation habitat (e.g., riparian scrub vs. cottonwood riparian
39 forest), and ability of habitat to support special-status species. DWR will

3.0 Environmental Setting, Impacts, and Mitigation Measures
3.5 Biological Resources—Aquatic

1 track habitat compensation efforts and only authorize implementation of
2 vegetation removal under the VMS at a rate and in locations consistent
3 with the volume and type of compensation habitat that has been
4 established. This habitat compensation tracking program will be included
5 in the program MMRP prepared to support this PEIR.

6 The plan must, at a minimum, meet the following basic performance
7 standard:

- 8 • Authorized losses of habitat do not exceed the function and value of
9 available compensation habitat.

10 DWR will coordinate with USFWS and DFG during preparation and
11 implementation of the plan to incorporate into the plan appropriate
12 compensation for effects on special-status species from vegetation
13 management along the levee system.

14 Various mechanisms may be employed to provide compensation habitat
15 under the plan, as long as the performance standard identified above is met.
16 The mechanisms include but are not limited to the following:

- 17 • Implementation of the CVFPP Conservation Strategy Framework
- 18 • Participation in existing NCCPs, HCPs, or other conservation plans
- 19 • Purchase of habitat credits at an established mitigation bank
- 20 • Habitat restoration implemented by a levee maintenance agency or
21 other entity

22 Any mitigation plantings in the floodway will not be permitted if they
23 would result in substantial increases in flood stage elevations, or alter flows
24 in a manner that would have a substantial adverse effect on the opposite
25 bank.

26 In many cases, implementing Mitigation Measures BIO-A-2a (NTMA),
27 and BIO-A-2b (NTMA) related to implementation of the VMS would
28 reduce impacts to an overall less-than-significant level and even sometimes
29 to a beneficial level. The extent, type, function, and values of any riparian
30 habitat removed would be fully compensated for by enhancing, restoring,
31 or creating riparian habitat elsewhere. However, removing riparian habitat
32 in some locations and enhancing, restoring, or creating habitat elsewhere
33 would result in overall relocation of riparian habitat within the Extended
34 SPA. It is possible that although some stream or river reaches may benefit
35 from compensatory habitat, habitat values in other stream or river reaches

1 could be substantially reduced, adversely affecting special-status fish
2 species that must move through these river reaches. Potential adverse
3 effects include increased predation risk, increased water temperatures, and
4 reduced food availability. In addition, planting vegetation in the floodway
5 may not be authorized by the Board, USACE, or other agencies if the
6 vegetation would impede flood flows sufficiently that a rise in water
7 surface elevation would cause a significant increase in risk to public safety.
8 Therefore, it cannot be assured that in all instances fisheries impacts would
9 be mitigated to a less-than-significant level. Therefore, Impact BIO-A-2
10 (NTMA) would be **potentially significant and unavoidable**.

11 **Impact BIO-A-3 (NTMA): *Effects on Special-Status Fish, Fish***
12 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
13 ***Critical Habitat, and Essential Fish Habitat Caused by Loss of Overhead***
14 ***Cover and Instream Woody Material during Construction***

15 Riverbank construction work for NTMAs could require not only removal
16 of overhead cover, but also removal from the river channel of IWM—an
17 important component of shaded riverine aquatic habitat. The loss of IWM
18 results in the loss of refugia for special-status fish from predators and high
19 flows. Riverbank construction could also reduce the amount of pool-
20 forming structures, and reduce the river channel's storage capacity for
21 sediment and organic matter as flows are passed more quickly downstream
22 (USFWS and NMFS1998). Note that material that can function as IWM
23 and vegetation that can provide shaded riverine aquatic habitat may be
24 above normal water levels in a river or stream system; thus, this material
25 does not provide habitat values at all times. However, when water
26 elevations rise, the IWM and vegetation becomes inundated or is near the
27 water surface, and it provides habitat benefits at these times.

28 IWM is particularly important to healthy riverine ecosystems, and may be
29 the most important structural component promoting stable fisheries
30 resources. Because IWM has a key role in maintaining the complexity of
31 essential habitat and refugia for special-status fish, the potential loss of
32 IWM could reduce both habitat quality and carrying capacity.

33 Riparian habitat provides structure (through shaded riverine aquatic
34 habitat) and food for fish. Shade decreases water temperatures, and low
35 overhanging branches can provide sources of food by attracting terrestrial
36 insects. As riparian areas mature and banks erode, the vegetation sloughs
37 off into the rivers. This process creates structurally complex habitat
38 consisting of IWM that offers refugia from predators, decreases water
39 velocities, and provides habitat for aquatic invertebrates. For these reasons,
40 many fish species are attracted to shaded riverine aquatic habitat,
41 particularly emigrating juvenile anadromous salmonids.

1 When riparian vegetation and soft substrates are lost, less organic material
2 (leaves, detritus, woody debris) enters the stream's ecosystem, which can
3 affect biological production at all trophic levels. The magnitude of these
4 effects depends on the degree to which riparian vegetation and natural
5 substrates are preserved or recovered during the life of the project.

6 Because overhead cover and IWM (and thus shaded riverine aquatic
7 habitat) would be lost as a result of construction along the banks and
8 levees, this impact would be **significant**.

9 **Mitigation Measure BIO-A-3 (NTMA): *Inventory and Replace Shaded***
10 ***Riverine Aquatic Habitat***

11 The project proponent will require that the following measures be
12 implemented to reduce the effects of program construction activities on
13 special-status fish, fish movement, nursery sites, riparian habitat,
14 designated critical habitat, and EFH. These measures may already be
15 incorporated into the conditions of permits identified above in Mitigation
16 Measure BIO-A-2a.

- 17 • An inventory of shaded riverine aquatic habitat will be conducted
18 before construction activities begin. Any shaded riverine aquatic habitat
19 that is removed will be replaced, with replacement to occur on site
20 when feasible. This includes IWM and other instream structures,
21 overhead shade, and shallow-water habitat.
- 22 • Mitigation credits may be purchased from a public or private mitigation
23 bank approved by USFWS and/or NMFS. The final number of credits
24 to be purchased will be determined by agency staff.
- 25 • A mitigation and monitoring plan will be developed and implemented
26 to ensure that the proposed bank treatments and any off-site mitigation
27 treatments fully compensate for losses of shaded riverine aquatic
28 habitat.

29 On-site revegetation is the preferred method of compensation, and could
30 reduce the impact to a less-than-significant level, and even potentially to a
31 beneficial level. If on-site compensation is not feasible, off-site mitigation
32 will be established either before or as soon as feasible after existing
33 vegetation is removed, or mitigation bank credits will be purchased before
34 existing vegetation is removed. As much of the mitigation habitat as
35 feasible will be created at or near the project site. If off-site mitigation is
36 necessary, a location that does not currently support riparian vegetation and
37 is capable of supporting riparian habitats will be preferred. Revegetation
38 requirements may be accomplished as part of implementation of the

1 CVFPP Conservation Framework. Any mitigation plantings in the
2 floodway will not be permitted if they would result in increases in flood
3 stage elevations, or alter flows affecting the opposite bank.

4 However, as described above at the conclusion of Mitigation Measure Bio-
5 A-2b, removing riparian habitat in some locations and enhancing, restoring,
6 or creating habitat elsewhere would result in overall relocation of riparian
7 habitat within the Extended SPA. It is possible that although some stream
8 or river reaches may benefit from compensatory habitat, habitat values in
9 other stream or river reaches could be substantially reduced, adversely
10 affecting special-status fish species that must move through these river
11 reaches. In addition, planting vegetation in the floodway may not be
12 authorized by the Board, USACE, or other agencies if the vegetation would
13 impede flood flows sufficiently that a rise in water surface elevation would
14 cause a significant increase in risk to public safety. Therefore, it cannot be
15 assured that in all instances fisheries impacts would be mitigated to a less-
16 than-significant level. Therefore, Impact BIO-A-3 (NTMA) would be
17 **significant and unavoidable.**

18 **Impact BIO-A-4 (NTMA): *Effects on Special-Status Fish, Fish***
19 ***Movement, Nursery Ground Usage, Designated Critical Habitat, and***
20 ***Essential Fish Habitat Caused by an Increase in Hydrostatic Pressure,***
21 ***Underwater Noise, and Vibrations during Construction***

22 Should any in-river construction work be needed for NTMAs, pile driving
23 could occur when cofferdams are placed to separate construction activities
24 from the active flowing channel. Pile-driving equipment and activities
25 would produce pressure waves and would create underwater noise and
26 vibration, thereby temporarily altering in-river conditions.

27 Hydrostatic pressure waves and vibration generated by pile driving can
28 adversely affect all life stages of fish. Effects on fish from changes in
29 hydrostatic pressure are not related to the distance of the fish from the point
30 of impact, but to the level and duration of the sound exposure (Hastings
31 and Popper 2005). Hydrostatic pressure waves may rupture the swim
32 bladders and other internal organs of all life stages of fish, and could
33 permanently injure their inner ears and lateral line organs (Caltrans 2001;
34 Hastings and Popper 2005). These injuries could reduce the ability of fish
35 (including special-status fish species) to orient in the water column, capture
36 prey, and reduce the ability of fish to avoid predators (Caltrans 2001).
37 Therefore, this impact would be **potentially significant.**

38 **Mitigation Measure BIO-A-4 (NTMA): *Conform to NMFS Guidelines***
39 ***for Pile-Driving Activities***

1 Several measures may be effective in reducing potential impacts on listed
2 fish species, either by decreasing the level of underwater sound or by
3 decreasing the number of fish exposed to the sound. The project proponent
4 and construction contractors will implement the following measures to the
5 extent feasible, as construction activities and site-specific conditions allow:

- 6 • Use fewer piles, smaller piles, or a different type of pile to minimize the
7 number and/or intensity of pile hammer impacts.
- 8 • Drive piles when species of concern are not present, as determined
9 either from surveys or by known migration and use patterns for species
10 occurring in the project area.
- 11 • Use a vibratory hammer rather than an impact hammer.
- 12 • Use a cushioning block between the hammer and pile.
- 13 • Use a confined or unconfined air bubble curtain.
- 14 • Drive piles during periods of reduced currents.

15 Pile-driving activities at project sites will be monitored to ensure that the
16 effects of pile driving on listed fish species are minimized. If any injury or
17 mortality to fish is observed, DFG, NMFS and/or USFWS will be
18 immediately notified and in-water pile driving will cease.

19 Implementing this mitigation measure would reduce Impact BIO-A-4
20 (NTMA) to a **less-than-significant** level.

21 **Impact BIO-A-5 (NTMA): *Effects on Special-Status Fish, Fish***
22 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
23 ***Critical Habitat, and Essential Fish Habitat Caused by Rock Placement***

24 Levee projects under the proposed program may involve placing rock
25 riprap material, generally on the waterside of the levee. Using riprap in
26 rivers or on the waterside of levee banks has been shown to affect natural
27 river processes and functions in all of the following ways:

- 28 • Reducing recruitment of spawning gravels
- 29 • Preventing new accretion of point bars and other deposition areas
30 where riparian vegetation can colonize
- 31 • Preventing meander migration, which over time reduces habitat
32 renewal, diversity, and complexity

- 1 • Limiting the channel's lateral mobility, potentially reducing habitat
2 complexity
- 3 • Decreasing nearshore roughness, thus causing water velocity to
4 increase at a high rate as discharge increases, which in turn may
5 accelerate erosion of earthen banks downstream
- 6 • Reducing the contribution of nutrient inputs to the stream by inhibiting
7 plant growth adjacent to the stream
- 8 • Reducing recruitment of IWM to the stream system
- 9 • Reducing benthic habitat, thus resulting in reduced abundance and
10 diversity of benthic macroinvertebrates

11 Protecting levee slopes with riprap generally results in nearshore hydraulic
12 conditions that are characterized by greater depths and faster, more
13 homogeneous water velocities than are found along natural banks. Higher
14 water velocities minimize deposition and retention of sediment and woody
15 debris. These changes reduce habitat complexity relative to habitat found
16 along natural shorelines, especially by eliminating the shallow, slow-
17 velocity habitat preferred by juvenile salmonids.

18 Replacing natural bank substrates with riprap can adversely affect
19 important ecosystem functions. Living space and food for terrestrial and
20 aquatic invertebrates is lost, eliminating an important food source for
21 special-status fish species. Part of the proposed program could involve
22 removing riprap and creating setback levees and floodplain habitat, which
23 would help offset the effects of placing any new levee riprap. In addition,
24 under the proposed program, vegetation could be incorporated into the rock
25 material of new and existing riprap, minimizing adverse effects. However,
26 a net increase in the extent of rock riprap on the SPFC could occur;
27 therefore, this impact would be **potentially significant**.

28 **Mitigation Measure BIO-A-5 (NTMA): *Implement Mitigation Measures***
29 ***BIO-A-2a and BIO-A-2b (NTMA)***

30 Mitigation Measures BIO-A-2a and BIO-A-3 include activities that would
31 minimize and compensate for adverse effects of rock placement on aquatic
32 resources. Additional opportunities may exist for on-site vegetation
33 planting as part of rock placement projects.

34 In many instances, implementing these mitigation measures could reduce
35 impacts to an overall less-than-significant level and even sometimes result
36 in a benefit to aquatic resources. However, replacing all vegetation and

1 IWM (and the resulting shaded riverine aquatic habitat) may not be
2 possible in all instances because some areas, especially urban areas, may
3 lack the right-of-way needed to implement vegetation replacement. In
4 addition, the planting of vegetation in the floodway may not be authorized
5 by the Board, USACE, or other agencies if the vegetation would impede
6 flood flows sufficiently that a rise in water surface elevation would cause a
7 significant increase in risk to public safety. Because it cannot be assured
8 that Impact BIO-A-5 (NTMA) can always be reduced to a less-than-
9 significant level, this impact would be **significant and unavoidable**.

10 **Impact BIO-A-6 (NTMA): *Effects on Special-Status Fish, Fish***
11 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
12 ***Critical Habitat, and Essential Fish Habitat Caused by the Increased***
13 ***Availability of Floodplain Habitat Generated by Setback Levees***

14 Numerous studies have found that floodplain habitat is valuable to native
15 fish species in the Central Valley. Seasonally flooded habitat provides
16 spawning, rearing, and foraging habitat for splittail and rearing habitat for
17 Chinook salmon (Sommer et al. 1997; Sommer et al 2001; Sommer et al.
18 2002; Baxter et al. 1996; Moyle et al. 2000; Jones & Stokes 1999).
19 Floodplain inundation benefits the fisheries by increasing habitat
20 availability and food supply and reducing predation rates. The duration and
21 timing of inundation are key factors in the success of splittail spawning and
22 rearing. A positive correlation exists between the number of days of
23 inundation and the abundance of juvenile splittail in years when floodplains
24 are inundated continuously for at least 4 weeks between March and April
25 (Sommer et al. 1997; Moyle et al. 2000; Jones & Stokes 2001).

26 Chinook salmon that rear in seasonally flooded habitat have higher survival
27 and growth rates than juveniles that remain in the main river channel to rear
28 (Jones & Stokes 1999; Sommer et al. 2001). When they rear in flooded
29 habitat, juvenile salmon have been found to have growth rates of more than
30 1 millimeter per day and as much as 20 millimeters in 2–3 weeks (Jones &
31 Stokes 2001). The water temperature is typically higher in floodplain
32 habitat than in habitats in main channels. Although increased temperature
33 increases metabolic requirements, the productivity in flooded habitat also
34 increases, resulting in higher growth rates (Sommer et al. 2001). The
35 production of drift invertebrates in the Yolo Bypass has been found to be
36 one to two times greater than in the Sacramento River (Sommer et al.
37 2001).

38 In addition, flooded vegetation support invertebrates that are a substantial
39 source of food for rearing juveniles. An increase in the areas of flooded
40 habitat can reduce the competition for food and space and potentially

1 reduce the possibility that these fish will encounter predators (Sommer et
2 al. 2001).

3 Setting back levees provides an opportunity to increase the amount of
4 floodplain habitat. Where setbacks expand a watercourse's floodway, this
5 new area available for inundation could provide additional resting and
6 rearing habitat for native fish species. Although expanded inundation areas
7 resulting from setback levees are not of the same magnitude as flooded
8 bypasses, these areas could contribute to habitat values, and they provide
9 an opportunity for development of shaded riverine aquatic and IWM
10 habitat that benefits these fish species.

11 However, stranding has been identified as a risk for juvenile fish that use
12 floodplain habitat, and this issue may occur in setback levee areas as well.
13 Fish stranding is a function of fish presence, hydrology, and topography.
14 Often, human-created structures such as borrow pits, artificial ponds, and
15 drainage ditches have resulted in fish stranding (Jones & Stokes 1999).
16 Typically, as flows recede, juvenile fish will move from the draining area.
17 Juvenile splittail have been observed moving off of the floodplain habitat
18 with receding flow (Sommer et al. 1997; Baxter et al. 1996). However,
19 juvenile Chinook salmon may not return to the main channel with receding
20 flows, resulting in stranding. Deeper ponds that have no outlet back to the
21 river can trap young fish, resulting in mortality if flows do not increase and
22 reestablish a connection with the river. Mortality may result when water
23 temperatures increase or oxygen levels decline while the water body
24 evaporates, when the water body evaporates completely, or from predation
25 by piscivorous (fish-eating) birds and mammals attracted to the trapped
26 fish. The ability of fish to return to the main channel appears to be
27 determined by size and readiness to migrate downstream in response to
28 smoltification (the physiological change allowing salmon to function in salt
29 water) (Jones & Stokes 1999).

30 Although creating new floodplain habitat can provide a substantial benefit
31 to fish populations, fish stranding in floodplain habitat created by a new
32 setback levee could outweigh these benefits through direct mortality to
33 special-status fish species. Therefore, this impact would be **potentially**
34 **significant**.

35 **Mitigation Measure BIO-A-6 (NTMA): *Design and Implement***
36 ***Floodplain Habitat to Minimize Stranding***

37 To avoid or minimize the potential for fish stranding associated with the
38 creation of new floodplain habitat, the existing topographic and hydrologic
39 characteristics of the floodplain will be examined to define the flooding

1 regime, drainage patterns, water depths, and potential risks of fish
2 stranding.

3 Potential floodplain habitat will slope to a main channel or slough to
4 facilitate complete drainage and avoid depressions or other low-lying
5 floodplain features that may strand fish. Periodic recontouring (e.g., filling
6 and excavation) of floodplain surfaces may be required to avoid stranding
7 fish.

8 Implementing this mitigation measure would reduce Impact BIO-A-6
9 (NTMA) to a **less-than-significant** level and assist in maximizing the
10 overall benefits of creating new floodplain habitat.

11 **3.5.5 Environmental Impacts and Mitigation Measures** 12 **for LTMA**s

13 This section describes the physical effects of LTMA

s on aquatic biological
14 resources. LTMAs include a continuation of activities described as part of
15 the NTMAs and all other actions included in the proposed program, and
16 consist of all of the following types of activities:

- 17 • Widening floodways (through setback levees and/or purchase of
18 easements)
- 19 • Constructing weirs and bypasses
- 20 • Constructing new levees
- 21 • Changing operation of existing reservoirs
- 22 • Achieving protection of urban areas from a flood event with 0.5 percent
23 risk of occurrence
- 24 • Changing policies, guidance, standards, and institutional structures
- 25 • Implementing additional and ongoing conservation elements

26 Actions included in the LTMA

s are described in more detail in Section 2.4,
27 “Proposed Management Activities.”

28 Impacts and mitigation measures identified above for NTMA

s would also
29 be applicable to many LTMAs and are identified below. The NTMA
30 impact discussions and mitigation measures are modified or expanded
31 where appropriate, or new impacts and mitigation measures are included if
32 needed, to address conditions unique to LTMAs. The same approach to
33 future implementation of mitigation measures described above for NTMAs

1 and the use of the term “project proponent” to identify the entity
2 responsible for implementing mitigation measures also apply to LTMA.

3 In addition, as described previously and in Section 3.1.2, “Analysis
4 Methodology,” because many LTMA are more general and conceptual,
5 additional impacts of those LTMA are also described below in a broader
6 narrative format, along with a list of suggested mitigation strategies that
7 could be applied to these impacts. This more general analysis is provided in
8 the subsection titled “LTMA Impact Discussions and Mitigation
9 Strategies.”

10 ***LTMA Impacts and Mitigation Measures***

11 ***Impact BIO-A-1 (LTMA): Potential Effects on Special-Status Fish,***
12 ***Fish Movement, Nursery Ground Usage, Riparian Habitat, Designated***
13 ***Critical Habitat, and Essential Fish Habitat Caused by Degradation of***
14 ***Water Quality during Construction or Operations and Maintenance***
15 ***Activities***

16 This impact would be similar to Impact BIO-A-1 (NTMA) because the
17 same impact mechanisms would occur. LTMA could occur throughout the
18 study area and could be larger in scale than NTMA; as a result, this impact
19 has a greater potential to occur than Impact BIO-A-1 (NTMA). However,
20 as described previously for NTMA, protective measures such as
21 implementation of a SWPPP are part of the LTMA, and compliance with
22 federal and State permits for construction activities would require
23 additional protective measures. Therefore, this impact would be **less than**
24 **significant**. No mitigation is required.

25 ***Impact BIO-A-2 (LTMA): Effects on Special-Status Fish, Fish***
26 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
27 ***Critical Habitat, and Essential Fish Habitat Caused by Loss of Overhead***
28 ***Cover and Instream Woody Material as Part of the Vegetation***
29 ***Management Strategy***

30 This impact would be similar to Impact BIO-A-2 (NTMA) because the
31 same impact mechanisms would occur. Therefore, this impact would be
32 **potentially significant**.

33 ***Mitigation Measure BIO-A-2 (LTMA): Implement Mitigation Measures***
34 ***BIO-A-2a (NTMA) and BIO-A-2b (NTMA)***

35 Implementing this mitigation measure would reduce Impact BIO-A-2
36 (LTMA), but it cannot be assured that the impact would be reduced to a
37 less-than-significant level in all cases. Impact BIO-A-2 (LTMA) would be
38 **potentially significant and unavoidable**.

1 **Impact BIO-A-3 (LTMA): *Effects on Special-Status Fish, Fish***
2 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
3 ***Critical Habitat, and Essential Fish Habitat Caused during Construction***

4 This impact would be similar to Impact BIO-A-3 (NTMA) because the
5 same impact mechanisms would occur. Therefore, this impact would be
6 **significant**.

7 **Mitigation Measure BIO-A-3 (LTMA): *Implement Mitigation Measures***
8 ***BIO-A-3 (NTMA)***

9 Implementing this mitigation measure would reduce Impact BIO-A-3
10 (LTMA), but it cannot be assured that the impact would be reduced to a
11 less-than-significant level in all cases. Impact BIO-A-3 (LTMA) would be
12 **significant and unavoidable**.

13 **Impact BIO-A-4 (LTMA): *Effects on Special-Status Fish, Fish***
14 ***Movement, Nursery Ground Usage, Designated Critical Habitat, and***
15 ***Essential Fish Habitat Caused by an Increase in Hydrostatic Pressure,***
16 ***Underwater Noise, and Vibrations during Construction***

17 This impact would be similar to Impact BIO-A-4 (NTMA) because the
18 same impact mechanisms would occur. Therefore, this impact would be
19 **potentially significant**.

20 **Mitigation Measure BIO-A-4 (LTMA): *Implement Mitigation Measure***
21 ***BIO-A-3 (NTMA)***

22 Implementing this mitigation measure would reduce Impact BIO-A-4 to a
23 **less-than-significant** level.

24 **Impact BIO-A-5 (LTMA): *Effects on Special-Status Fish, Fish***
25 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
26 ***Critical Habitat, and Essential Fish Habitat Caused by Rock Placement***

27 This impact would be similar to Impact BIO-A-5 (NTMA) because the
28 same impact mechanisms would occur. Therefore, this impact would be
29 **potentially significant**.

30 **Mitigation Measure BIO-A-5 (LTMA): *Implement Mitigation Measure***
31 ***BIO-A-5 (NTMA)***

32 Implementing this mitigation measure would reduce Impact BIO-A-4
33 (LTMA), but not to a less-than-significant level. Impact BIO-A-4 (LTMA)
34 would be **potentially significant and unavoidable**.

1 **Impact BIO-A-6 (LTMA): *Effects on Special-Status Fish, Fish***
2 ***Movement, Nursery Ground Usage, Riparian Habitat, Designated***
3 ***Critical Habitat, and Essential Fish Habitat Caused by the Increased***
4 ***Connectivity of Floodplain Habitat Generated by Setback Levees***

5 This impact would be similar to Impact BIO-A-6 (NTMA) because the
6 same impact mechanisms would occur. Therefore, this impact would be
7 **potentially significant**.

8 **Mitigation Measure BIO-A-6 (LTMA): *Implement Mitigation Measure***
9 ***BIO-A-6 (NTMA)***

10 Implementing this mitigation measure would reduce Impact BIO-A-6
11 (NTMA) to a **less-than-significant** level and assist in maximizing the
12 overall benefits of creating new floodplain habitat.

13 **Impact BIO-A-7 (LTMA): *Effects on Passage by Special-Status Fish***
14 ***and Fish Movement***

15 Structural modifications at weirs would help improve fish passage in ways
16 that would reduce migration delays and reduce predation. This improved
17 passage would be beneficial to special-status fish species and fish
18 movement. This impact would be **beneficial**. No mitigation is required.

19 ***LTMA Impact Discussions and Mitigation Strategies***

20 Because of the more general and conceptual nature of many LTMA's, a
21 great deal of uncertainty exists about how some LTMA's may be
22 implemented and what environmental effects may result from their
23 implementation. This uncertainty is to be expected for a broad, multiyear,
24 and in some areas, conceptual program such as the CVFPP. Although these
25 uncertainties exist, sufficient information exists to at least disclose
26 additional potential impacts of LTMA's besides those discussed in the
27 impact/mitigation pairings above. The following additional LTMA impacts
28 are described in a broad narrative format; because of the uncertainty
29 surrounding these impacts, no determination regarding their significance is
30 provided. Consistent with Section 15145 of the CEQA Guidelines, these
31 impacts are too speculative for evaluation beyond the narrative disclosure
32 provided here.

33 Future project-specific CEQA evaluations for individual LTMA's will be
34 used to determine the potential for the impacts described below to occur,
35 determine their level of significance, and identify project-specific
36 mitigation measures for significant impacts. Examples of potential
37 mitigation strategies are provided after the following narrative impact

1 discussions to disclose the nature and extent of mitigation actions that
2 might be necessary to address these impacts.

3 For more information on this approach to evaluating LTMA impacts and
4 providing mitigation strategies, see Section 3.1.2, “Analysis Methodology.”

5 Impact discussions are divided among the geographic areas in the program
6 study area (i.e., Extended SPA, Sacramento and San Joaquin Valley
7 watersheds, and SoCal/coastal CVP/SWP service areas). They are further
8 subdivided according to the type of action (i.e., construction of conveyance
9 facilities, facilities operations and maintenance from storage or conveyance
10 actions, and other management actions).

11 **LTMA Impact Discussions**

12 *Extended Systemwide Planning Area*

13 *Construction of Conveyance Facilities* Construction-related
14 impacts of LTMA on aquatic biological resources are thoroughly
15 described and evaluated above in the analysis of NTMA and LTMA. A
16 more general narrative description of additional construction-related
17 LTMA impacts in the Extended SPA is not required.

18 *Facilities Operations and Maintenance from Storage or*
19 *Conveyance Actions* Operational changes in multiple Central Valley
20 reservoirs as would occur under the LTMA may affect the aquatic habitat
21 conditions of downstream rivers. The full extent of potential operational
22 changes and the possible interactions of those changes in multiple
23 reservoirs are unknown. As a result, altered flow patterns could benefit fish
24 in some instances and adversely affect them in other instances. Generally,
25 changing reservoir operations would minimize damaging peak flood flow
26 releases. Such changes could include using forecast-based operations
27 (increasing reservoir releases earlier than would otherwise occur in
28 anticipation of incoming storms) or coordinating reservoir operations to
29 minimize peak downstream flows. These actions, however, could also
30 slightly reduce the amount of water available for later releases if the
31 reservoir releases were not fully offset by inflows from storms. The
32 following beneficial and adverse changes to important river dynamic
33 processes could occur:

- 34 • Redd scouring or siltation
- 35 • Alteration of ecologically important geomorphic processes resulting
36 from altered frequency and magnitude of high flows (channel forming
37 and maintenance)

- 1 • Altered floodplain inundation

2 Because the locations and extent of proposed changes in reservoir
3 operations have not been defined, any effects on fish resulting from flow
4 changes associated with LTMA are speculative at best, and the impact
5 could range from potentially significant to beneficial.

6 Changes in flow and reservoir operations could also affect water
7 temperatures in Central Valley rivers. Fish species have different responses
8 to water temperature conditions depending on their physiological
9 adaptations. In general, salmonids have evolved under conditions where
10 water temperatures are fairly cool. Delta smelt and splittail can tolerate
11 warmer temperatures. In addition to species-specific thresholds, different
12 life stages have different water temperature requirements. Eggs and larval
13 fish are the most sensitive to warm water temperatures.

14 Because the location and extent of proposed changes in reservoir
15 operations have not been defined, any effects on fish resulting from
16 changes in water temperature associated with LTMA are speculative at
17 best, and the impact could range from potentially significant to beneficial.

18 Fluctuations in reservoir elevations could affect the spawning and rearing
19 of reservoir fishes, particularly bass species. Water-level fluctuations
20 associated with reservoir management are perhaps the most important
21 factor affecting reservoir fish. Water-level fluctuations directly affect black
22 bass species, such as largemouth bass (*Micropterus salmoides*) and spotted
23 bass (*M. punctulatus*), which construct nests for their eggs in shallow-water
24 habitat (Kohler et al. 1993; Thorton et al. 1990; Aasen and Henry 1980).
25 Falling water levels expose nests to wave action or dewater them entirely,
26 while rising water levels may expose the nests to cold water that kills the
27 eggs or slows their development.

28 Because proposed changes in operations have not been defined, any effects
29 on fish resulting from changes in reservoir elevation fluctuations associated
30 with LTMA are speculative at best, and the impact could range from
31 potentially significant to less than significant.

32 *Other Management Actions* Impacts on aquatic biological
33 resources resulting from “other management actions” included in LTMA
34 are thoroughly described and evaluated above in the analysis of NTMA
35 and LTMA. A general narrative description of additional LTMA impacts
36 related to other management actions in the Extended SPA is not required.

1 *Sacramento and San Joaquin Valley Watersheds*

2 *Facilities Operations and Maintenance from Storage or*
3 *Conveyance Actions* The mechanisms for and effects of operating and
4 maintaining storage facilities in the Sacramento and San Joaquin Valley
5 watersheds would be similar to those described above for operating and
6 maintaining water storage facilities in the Extended SPA. Mitigating these
7 potentially significant adverse impacts to less-than-significant levels may
8 not always be possible.

9 None of the program’s management actions related to conveyance would
10 be implemented in the portions of the Sacramento and San Joaquin Valley
11 watersheds outside the Extended SPA. Therefore, no impacts on aquatic
12 biological resources would result from conveyance-related management
13 actions in this area.

14 *Other Management Actions* Impacts on aquatic biological
15 resources of “other management actions” included in the LTMA are
16 thoroughly described and evaluated above in the analysis of NTMA and
17 LTMA. A general narrative description of additional LTMA impacts
18 related to other management actions in the Sacramento and San Joaquin
19 Valley watersheds is not required.

20 *SoCal/Coastal CVP/SWP Service Areas* None of the program’s
21 management actions would be implemented in the SoCal/coastal
22 CVP/SWP service areas. There would also be no substantial or long-term
23 changes to water deliveries in this area and reservoir levels would be
24 affected only minimally, if at all. Therefore, no impacts on aquatic
25 resources would occur.

26 **LTMA Mitigation Strategies** The following mitigation strategies are
27 example of approaches that may be considered to address significant
28 impacts via the mechanisms described above. These mitigation strategies
29 may be considered, as applicable, during project-level evaluation of
30 specific LTMA. For more information on LTMA mitigation strategies, see
31 Section 3.1.2, “Analysis Methodology.”

32 Specific mitigation measures identified above in the NTMA and LTMA
33 impact/mitigation pairings are not identified again in the mitigation
34 strategies. It is assumed that mitigation measures described in the
35 impact/mitigation pairings above would already be required, as applicable,
36 as part of the project-level evaluation of specific LTMA. Not all
37 mitigation strategies will apply to all LTMA; the applicability of
38 mitigation strategies will vary based on the location, timing, and nature of
39 each management action. In addition, some mitigation strategies on their
40 own may not constitute sufficient mitigation under CEQA but must be

1 coupled with other mitigation strategies to reduce the impacts of LTMA's to
2 less-than-significant levels.

3 The following potential mitigation strategies have been identified for
4 aquatic biological resources, if necessary and feasible, to minimize certain
5 types of significant or potentially significant impacts from project-level
6 LTMA's:

- 7 • Create or enhance in-reservoir habitat in the form of woody debris or
8 other hard structures to provide refugia and/or shallow-water spawning
9 habitat for black bass or other fish species of high recreational value in
10 reoperated reservoirs.
- 11 • Maintain or increase cold-water storage in reservoirs to minimize
12 temperatures in downstream rivers below reoperated reservoirs.
- 13 • Minimize reservoir releases that increase downstream flow fluctuations
14 and adversely affect the success of salmon spawning below reoperated
15 reservoirs.
- 16 • Ensure that any changes to reservoir operations continue to allow
17 sufficient flow releases downstream to meet all relevant minimum and
18 other instream flow requirements below reoperated reservoirs.
- 19 • Ensure that any changes to reservoir operations maintain downstream
20 riverine geomorphic conditions that maintain or enhance physical
21 habitat conditions for all runs of salmon, steelhead, and other special-
22 status fish species below reoperated reservoirs.

23